When modelled debris arrives at each find site, compared with the date it was found



Plain English. Each row is one of the nine confirmed or almost-certain debris items that has a reliable find date. Thick bar: the middle half of the modelled arrival dates at that site; thin line: 90 % of arrival dates; white tick: median. The number at the right is the share of the modelled arrivals that came before the find date. The real item arrived before it was found, so low numbers mean the find was early for that start point. From the core, the three earliest finds (Réunion, Mossel Bay, Paindane) were early: only 15-23 % of modelled arrivals came first. From 30 S, 37-55 % came first. This agrees with the drift result, which prefers an impact north of the core. This check is not a likelihood; it does not include the chance that the debris was found and reported.

Technical. Snapshot harness on drift d51d80a (untracked), first-beaching day, refloat off, 15-160 E, two models equal weight (GLORYS12V1 0.494 m + ERA5; GlobCurrent 0 m + ERA5, windage U(-0.75,-0.60) % U10), classes pooled. Core: 200 draws from run J waypoints D0' (seeds 1-8, evidence-weighted), ~10,000 particles each per model; 30 S: 7th-arc point 97.775 E, 100,008 per model. Sites: primary items of debris-evidence-audit.csv (decision primary*), beach cells (0.5 deg) with centre within 100 km. Sum over finds of ln(share of released objects arriving there by the find date): core -52.4, 30 S -43.1, Pleiades objects -48.1, 40 S -59.6 (descriptive only).