




The 00:19 data options (bto-bfo)

Each option scores the same descents with a different choice of the 00:19 transmissions. Its log-evidence increment is how well the flight posterior, continued to impact, predicts those data: a strongly negative value means the descents cannot produce them. ESS is the effective number of descents (and of 00:11 trajectories) that carry the result.



Set core-bto: placeholder

Case bto-bfo; 00:19 data option both-bto; modules none. First water contact.



UNCONVERGED. Latitude median -37.77° , 95% interval -39.89° to -35.70° . Log evidence of the set's data under this hypothesis: -10.48 (replicates -10.48, -10.46). Split-half agreement: latitude 0.86, map 0.27.


Effective sample sizes (descents / trajectories): combined 17,292 / 1,747; option 17,292 / 1,747. Share of the pre-module posterior not computed: set 0.00%.

Alternatives, posterior and evidence per option: none.

Predictions (median [2.5%, 97.5%]): none.

Set core-bto: the average over end-of-flight hypotheses

The end-of-flight hypotheses cannot be ranked by a prior that data could test, so their average is shown only under two priors: the terminal module's own, and uniform. Each hypothesis's weight is its prior times the evidence of the set's data under it.





Prior terminal module: median -37.77°, 95% -39.89° to -35.70°; hypothesis weights placeholder 1.00.

Prior uniform: median -37.77°, 95% -39.89° to -35.70°; hypothesis weights placeholder 1.00.

Set core-all: placeholder

Case bto-bfo; 00:19 data option all; modules none. First water contact.



UNCONVERGED. Latitude median -37.64° , 95% interval -39.73° to -35.50° . Log evidence of the set's data under this hypothesis: -52.62 (replicates -52.63, -52.51). Split-half agreement: latitude 0.82, map 0.26.


Effective sample sizes (descents / trajectories): combined 16,550 / 1,668; option 16,550 / 1,668. Share of the pre-module posterior not computed: set 0.00%.

Alternatives, posterior and evidence per option: final-bfo-model: inflated 1.00 (ln Z -51.52), no-offset 0.00 (ln Z -607.35), startup-offset 0.00 (ln Z -708.79).

Predictions (median [2.5%, 97.5%]): none.

Set core-all: the average over end-of-flight hypotheses

The end-of-flight hypotheses cannot be ranked by a prior that data could test, so their average is shown only under two priors: the terminal module's own, and uniform. Each hypothesis's weight is its prior times the evidence of the set's data under it.




Prior terminal module: median -37.64°, 95% -39.73° to -35.50°; hypothesis weights placeholder 1.00.

Prior uniform: median -37.64°, 95% -39.73° to -35.50°; hypothesis weights placeholder 1.00.

Set plumbing: placeholder

Case bto-bfo; 00:19 data option r600-bto; modules arc-kernel. First water contact.



UNCONVERGED. Latitude median -37.81° , 95% interval -39.92° to -35.71° . Log evidence of the set's data under this hypothesis: -5.27 (replicates -5.27, -5.32). Split-half agreement: latitude 0.85, map 0.27.


Effective sample sizes (descents / trajectories): arc-kernel 24,896 / 2,490; combined 22,831 / 2,292; option 22,831 / 2,292. Share of the pre-module posterior not computed: arc-kernel 0.00%, set 0.00%.

Alternatives, posterior and evidence per option: none.

Predictions (median [2.5%, 97.5%]): none.

Set plumbing: the average over end-of-flight hypotheses

The end-of-flight hypotheses cannot be ranked by a prior that data could test, so their average is shown only under two priors: the terminal module's own, and uniform. Each hypothesis's weight is its prior times the evidence of the set's data under it.



Prior terminal module: median -37.81°, 95% -39.92° to -35.71°; hypothesis weights placeholder 1.00.

Prior uniform: median -37.81°, 95% -39.92° to -35.71°; hypothesis weights placeholder 1.00.