




Seabed search coverage: Phase 2 valid-data fraction (blue); OI 2018 outline less Phase 2, inferred (orange)




Phase 2 deep-tow: valid-pixel fraction per 0.01 deg cell




Phase 2 go-phoenix: valid-pixel fraction per 0.01 deg cell




Phase 2 dhj: valid-pixel fraction per 0.01 deg cell



Phase 2 auv: valid-pixel fraction per 0.01 deg cell





Area checks

Phase 2 deep-tow: valid 5 m pixels	103,921.5 km2	not seen by another sensor: 88,507 km2
Phase 2 go-phoenix: valid 5 m pixels	14,627.3 km2	not seen by another sensor: 11,901 km2
Phase 2 dhj: valid 5 m pixels	3,164.3 km2	not seen by another sensor: 2,303 km2
Phase 2 auv: valid 5 m pixels	16,164.1 km2	not seen by another sensor: 818 km2
Phase 2 union (per-cell maximum; used)	120,486.5 km2	per-cell sum bound 120,488 km2; ATSB ~120,000 km2
Phase 2 union, module raster (0.01 deg, 8-bit)	120,489.9 km2	
Deep-tow WFS footprint, spherical polygon area	103,924.7 km2	its attribute: 130,961 = square degrees x 111.12^2
Deep-tow WFS footprint, rasterised	103,922.0 km2	-0.00% vs polygon
Deep-tow pixels outside the WFS footprint	100.4 km2	
WFS footprint without deep-tow pixels	100.9 km2	
G0 Phoenix outside the deep-tow WFS footprint	13,445.7 km2	brief: ~14,400 km2
Bluefin-21 display polygons, spherical	771.4 km2	ATSB reports 860 km2
Bluefin-21, module raster	771.4 km2	
OI 2018 outline (grade C), spherical	148,993.2 km2	OI: >112,000 km2 (May 2018), 120,000 km2 (data donation)
OI 2018 outline over Phase 2 coverage	22,980.0 km2	
OI 2018 layer: outline less Phase 2, module raster	126,009.3 km2	reported area / layer: 0.889-0.952 (coverage_fraction)
PASS: deep-tow WFS layer vs the 31 Aug 2026 snapshot	(sha256 6591be13bbd3)	