
Where the aircraft hit the ocean, by latitude (core run I, not converged)



(i) The curves show how probable each impact latitude is. The data at 00:19 change the curves only a little. The seabed searches found nothing, so they move probability away from the searched area, to the north. The waypoint family has almost all of the weight because the four families have equal prior weight; the free-flight curve is shown beside it. The result is not converged. Dotted curves have too few effective samples.
(ii) Technical: Core run I: 4 core families x 8 seeds x 100,000 hand-off rows, equal family prior; NOT CONVERGED (split-half waypoints 0.916, floor 0.924; free flight 0.830, descent-climb 0.794, Davey reproduction 0.891). End of flight run I (binary mh370-eof-ri1, 102.4 M impacts): piloted turns and interim pitch-down provisional; no structural limit (G13). 00:19: competing log-on causes q = 1/3, W = 00:10:58-01:15:56, C2 silent, survival factor applied; no treatment uses the R1200 BTO. Ocean: GLORYS12 and GlobCurrent (product-relative windage), equal prior, mixed per row (Bayesian model averaging). Drift h50; unscored rows carried at the mean ratio (<= 0.11 % of weight); Western Australia non-recovery not modelled (G26). Hydroacoustics predictive only, not in these products. Searches: ATSB Phase 2 + Bluefin-21, rho 0.05, point target; Ocean Infinity excluded. Pléiades: rho 0.25, rating 5, six clusters plus four COSMO-SkyMed contacts, conditional on the objects being debris. Strata mixed by core share x 00:19 evidence (ruling C), conditional on equal family prior. Composer pass 2, architecture stand-in. Curves: 0.05° latitude bins, Gaussian 0.1°. (a, b) all six 00:19 treatments; (c) 00:19 R600 BTO Only; (d) R600 BTO Only + drift, per core family (waypoints 0.9982 of the equal-prior mixture).