Two ocean models for the debris drift, weighted by the evidence (core run I, not converged)





(i) Two ocean models give the drift of the debris. Each model starts with a probability of one half. The debris finds change this probability only a little, so the result uses both models together. With the 00:19 timing value and the drift, the two medians are 0.2° apart.
(ii) Technical: Core run I: 4 core families x 8 seeds x 100,000 hand-off rows, equal family prior; NOT CONVERGED (split-half waypoints 0.916, floor 0.924; free flight 0.830, descent-climb 0.794, Davey reproduction 0.891). End of flight run I (binary mh370-eof-ri1, 102.4 M impacts): piloted turns and interim pitch-down provisional; no structural limit (G13). 00:19: competing log-on causes q = 1/3, W = 00:10:58-01:15:56, C2 silent, survival factor applied; no treatment uses the R1200 BTO. Ocean: GLORYS12 and GlobCurrent (product-relative windage), equal prior, mixed per row (Bayesian model averaging). Drift h50; unscored rows carried at the mean ratio (<= 0.11 % of weight); Western Australia non-recovery not modelled (G26). Hydroacoustics predictive only, not in these products. Searches: ATSB Phase 2 + Bluefin-21, rho 0.05, point target; Ocean Infinity excluded. Pléiades: rho 0.25, rating 5, six clusters plus four COSMO-SkyMed contacts, conditional on the objects being debris. Strata mixed by core share x 00:19 evidence (ruling C), conditional on equal family prior. Composer pass 2, architecture stand-in. P(GLORYS12) = Z_G / (Z_G + Z_C) on the core mixture; the GlobCurrent run is partly in-sample for drifter skill, so no up-weighting. Treatments with both BFOs have low effective sample size after drift (see the coverage table).