
If the imaged objects are from 9M-MRO: where the debris entered the sea, on GEBCO 2026 relief — 00:19 R600 BTO — core run I, NOT CONVERGED

Pléiades + all four COSMO-SkyMed contacts, one debris field after Phase 2 + Bluefin-21 (official)



STE100: This map shows where the aircraft possibly went into the sea, if the objects in the Pléiades satellite images are from 9M-MRO. The red bands show the 50, 90 and 99 % regions with this assumption. The points show the probability without this assumption. The drawn result uses two ocean models with equal weight and a correlation of 0.25 between the drift errors of the two satellite data sets. Only the official searches (Phase 2 and Bluefin-21) are applied. The flight calculation (core run I) is not converged. Use these results with caution.

Technical: core run I (4 strata x 8 seeds; waypoints carries core P 0.9982) NOT CONVERGED; end-of-flight physics PROVISIONAL (piloted-turn model, INTERIM pitch-down, push-over and Boeing sequence; end of flight's run I labels). 00:19 R600 BTO; log-on cause: competing-causes total, q = 1/3, C2 silent (primary); survival factor applied; end of flight's own row weights (option_table_0019), strata by core P x own evidence. Search: searched areas' per-impact ln P(no find), rho 0.05, point target, Phase 2 (q 0.945) + Bluefin-21 (q 0.9); OI 2018 and 2025-26 not applied (grade C; drawn only). Under H: 12 rating-5 Pléiades objects (6 clusters, 3 km, equal weight) x all four COSMO-SkyMed contacts, one debris field; bivariate-normal transport errors, ρ = 0.25 (measured 0.07-0.41), ρ = 0 / 0.5 beside; GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight, GlobCurrent windage product-relative (drift audit F1). PROPOSED headline; GLORYS12 only (ruling 2) beside. Maps from per-impact columns on the wide 0.05 deg grid (prepare/runi_closeups.py). Conditional on H; no identity likelihood, no Bayes factor or P(H | data).

Coverage: effective impacts after the official searches 42,142 under H (372,244 without H); A1 under H 6,284. Gate (1,000) met.

Result, after the official searches: 90 % region 64,426 km², mean 35.56 S 92.22 E; tension ln S -1.07 (p 0.19), mean shift 104 NM; PDF without H inside the 90 % region 19 %. Beside it, GLORYS12 only at ρ = 0.25: 56,120 km², 35.61 S 92.25 E, ln S -0.91.

Relief: GEBCO 2026 15-arcsecond grid (doi:10.5285/4f68d5c7-45eb-f999-e063-7086abc036fa), shown at 0.025 deg; contours every 1 km depth. Right: probability per degree of latitude over the whole grid, after Phase 2 + Bluefin-21 (official).

Background: the impact PDF without H (flight posterior alone, after the same searches), as 6,000 points drawn in proportion to its probability from 0.05 deg grid cells and jittered within each cell. 60 % of it lies inside this frame.