
If the imaged objects are from 9M-MRO: where the debris entered the sea — 00:19 R600 BTO + both BFOs, Holland transient bias added (H1 da
core run I, drift-error correlation 0.25, two ocean models (proposed headline) — NOT CONVERGED — NOT ESTIMABLE (too few samples under H)



	ρ = 0.25 two models (drawn)	ρ = 0.25 GLORYS12 only	ρ = 0 two models	ρ = 0.5 two models
50 % region, km²	2,639	2,209	2,438	2,814
90 % region, km²	16,756	15,139	14,612	19,117
mean position	35.61 S 92.62 E	35.67 S 92.61 E	35.59 S 92.63 E	35.63 S 92.60 E
outside official searches	78.5 %	81.1 %	77.9 %	78.5 %
share left by searches	0.396	0.438	0.389	0.396
tension ln S (p)	-0.23 (0.20)	+0.04 (0.23)	-0.32 (0.21)	-0.16 (0.19)
mean shift from PDF without H	73 NM	71 NM	75 NM	72 NM
PDF without H inside 90 % region	18 %	17 %	16 %	21 %

STE100: This map shows where the aircraft possibly went into the sea, if the objects in the Pléiades satellite images are from 9M-MRO. The blue colour shows the probability with this assumption. The grey points show the probability without this assumption. The drawn result uses two ocean models with equal weight and a correlation of 0.25 between the drift errors of the two satellite data sets; the thin lines show the 90 % region for other settings. Only the official searches (Phase 2 and Bluefin-21) are applied. The flight calculation (core run I) is not converged. Use these results with caution. Technical: core run I (4 strata x 8 seeds; waypoints carries core P 0.9982) NOT CONVERGED; end-of-flight physics PROVISIONAL (piloted-turn model, INTERIM pitch-down, push-over and Boeing sequence; end of flight's run I labels). 00:19 R600 BTO + both BFOs, Holland transient bias added (H1 data); log-on cause: competing-causes total, q = 1/3, C2 silent (primary); survival factor applied; end of flight's own row weights (option_table_0019), strata by core P x own evidence. Search: searched areas' per-impact ln P(no find), rho 0.05, point target, Phase 2 (q 0.945) + Bluefin-21 (q 0.9); OI 2018 and 2025-26 not applied (grade C; drawn only). Under H: 12 rating-5 Pléiades objects (6 clusters, 3 km, equal weight) x all four COSMO-SkyMed contacts, one debris field; bivariate-normal transport errors, ρ = 0.25 (measured 0.07-0.41), ρ = 0 / 0.5 beside; GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight, GlobCurrent windage product-relative (drift audit F1). PROPOSED headline; GLORYS12 only (ruling 2) beside. Maps from per-impact columns on the wide 0.05 deg grid (prepare/runi_closeups.py). Conditional on H; no identity likelihood, no Bayes factor or P(H | data). Coverage: effective impacts after the official searches 460 under H (2,865 without H); A1 under H 124. NOT ESTIMABLE: below the 1,000 gate. The speckled regions, the small areas and the tension are sampling artefacts of too few effective impacts, not a measured result. Background: 6,000 points from the impact PDF without H after the same searches.