
With a continued dive in reach, the R1200 impact peak moves from 37.65° S to 37.35° S and the 90 % area shrinks to 47k km²



Note 1: Each panel shows the area that holds 50, 90 and 99 % of the probability of the ocean impact, for one way to use the two 00:19 satellite messages. The heavy black, brown and green lines show sea floor that was searched; the search result is not applied in these maps. The log-on cause mixes fuel exhaustion (weight 3) and other causes (weight 1). A piloted steep descent can now continue to the sea. The results are provisional.
Note 2: End of flight on core run I, 4 route families x 8 seeds x 100,000 hand-off particles at 00:11 (family weights 0.9982/0.0011/0.0004/0.0002); N = 16 children x 4 descents for waypoints, 8 for the others, binary mh370-eof-ri3: run I recipe + C2-silent aimed proposal + dive branch (each piloted push-over continues to the surface or levels off; U[0,1] hyper-prior, marginal 0.5) + SLS thrust 406.8 kN (EASA E.047). Competing causes C1 (Erlang lag) + C2 (1/W, W = 00:10:58-01:15:56 UTC, silent), q = 1/3; survival factor applied; no treatment uses the R1200 BTO. Mixed = equal-prior average of D3/D4/D5 by ln Z. Density 0.02° grid, Gaussian sigma 0.08°. Searched ground: coverage >= 0.5 outline (Bluefin-21 2014 not drawn). PROVISIONAL: piloted-turn model, INTERIM pitch-down. Core run I not converged (8-seed floor 0.916).