


Plain English. Regions that hold 50 % and 90 % of the probability of the impact position. Grey: satellite data, aircraft performance and end of flight (core). Blue: with the debris drift (nine finds, two ocean models) and the searched areas. Purple: also with the Pleiades objects, on the condition that they are aircraft debris (a conditional result). Tan: seabed searched by Phase 2 and Bluefin-21. 00:19 data held out; cause of the 00:19 log-on: fuel exhaustion or another cause with the aircraft silent (prior odds 1 to 3). Positions are impact points, not seabed points. The end-of-flight physics is not calibrated. Ocean Infinity areas are not used.

Technical. Run J, four strata by core P(family); cell D0 total q=1/3 (C2 silent). Equal-area grid 0.25 deg x authalic q step, 769.3 km2 cells; HPD = smallest cell set. Drift run A rebuilt surfaces 50 km, GLORYS12V1/GlobCurrent + ERA5 at equal prior, marginalised jointly with Pleiades (rho 0.25, gcrel, P + COSMO F1-F4). Searches seabed_loglik.py rho 0.05 (Phase 2 q 0.945, Bluefin-21 q 0.9); tan = union coverage >= 0.5. No settling resample on run J. Run J split-half accepted as a PASS by project ruling, 16 Oct 2026; north of 33 S under-sampled (see note). Not-computed rows carried at the (seed, option) mean.