


# Where the wreckage is on the seabed, for each use of the 00:19 satellite data

Flight families weighted by the flight model, then by the 00:19 data



Summary (STE): Grey areas show where the wreckage is on the seabed. Dashed orange lines show where the aircraft hit the water. The movement of the wreckage through the water adds less than 1 % to each grey area. The flight model is not yet stable, so these areas can change. There are not yet sufficient samples to show a result for Holland H1 and H2.

Technical: Settled-mass seabed density (real ocean: GLORYS12V1 column 75-115 E, 45-10 S, TEOS-10 density; GLORYS12V1 surface current; ERA5 wind; AusSeabed then GEBCO\_2026), settling 5b595bf, breakup table PROVISIONAL, afloat pieces excluded. Impacts: end of flight next-run on core (b) (4 strata x 4 seeds, prior track 289.7 deg, EoF 43262c31), option\_posteriors with +unpowered (airborne at 00:19:37, not powered at 01:15:56). Strata weighted by core's probability of each flight family, then by end of flight Z\_00:19 (+alive) for each family; P(family) for R600 BTO + Raw BFO = 0.639 / 0.140 / 0.204 / 0.018 (free / repro-radar / descent-climb / routes). Log-on cause: other (H1: fuel exhaustion). Systematic resample, one settling draw each: 200,000 (a), 40,000 (b-e). 0.02 deg grid, Gaussian 0.1 deg, HPD. Mixture ESS by Kish; below 1,000 not estimable (H1/H2: descent-model and sampler changes await Pete; architecture burst study). Labels: core (b) split-half NOT converged; two-tank bookkeeping only; internal-v1 one-engine flow 2x; dive class (b) and Boeing glide PROVISIONAL-OVERNIGHT; Mac, 2 threads.