


Conditional branch under H — 00:19 option: r1200/inflated



Source: end of flight's next-run impacts on core (b) m0011 hand-offs (EoF 3c6319f, idle floor ON), 4 strata mixed with core's P(family) held fixed and not updated by terminal-stage evidence (free 0.6948, Davey dynamics + radar 0.1527, descent-climb 0.1376, routes 0.0149; core split-half 0.71-0.88 vs floor 0.896, unconverged); within each stratum 4 seeds pooled with equal weight, 3.2 M impacts per seed. Free stratum detail: impacts next-free (4 seeds pooled with equal weight; source run /Users/pete/claude-science/orgs/9db41e8b-db54-4736-82b9-d77e2a9ad222/workspaces/23b26fb9-3c70-4c8e-909b-ac78b132214a/runs/src-next-free, code 3c6319f); prior track 289.7 deg (sd 1.0); config chain (run.json): davey2016-inmarsat + no-exhaustion-prior + s1-factor + s2-temperature + s3-internal + s4-ceiling + s5-hard-reject + reference-snapshots + radar-full + free + server-free + hypotheses/end-of-flight/smoke/snapshot-m0011.toml + hypotheses/end-of-flight/smoke/terminal.toml + hypotheses/end-of-flight/full/reference-289.toml + hypotheses/end-of-flight/full/descent-idle-floor.toml + hypotheses/end-of-flight/full/seed-1.toml; tempered sampler (m1839, m1941, m2041, m2141, m2241, m0011, 16 stages); particles per mode [500000, 250000, 1250000, 250000, 1250000]; BFO bias (run.json bfo_bias: {"mean_hz":150.0,"sd_hz":25.0}).

Labels: core (b) split-half NOT converged; two-tank bookkeeping only; PROVISIONAL-OVERNIGHT; EoF sweep run by an architecture stand-in on the Pleiades module's behalf, module to review; strata MIXED by P(family).

00:19 data option 'r1200/inflated' (project sensitivity): R1200 BFO scored with an independent zero-mean inflated BFO error; R600 held out; log-on cause 'other' (no fuel-exhaustion log-on density). Weight = impact weight x exp(loglik:r1200/inflated).

End-of-flight physics PROVISIONAL: dive class (b) (being rebuilt against Boeing's simulator runs); Boeing-calibrated glide (Pete's reference).

Search (row 2): the searched-areas module's own per-impact likelihood, base case Phase 2 2014-17 + Bluefin-21, point target, undetectable probability rho = 0.05; Ocean Infinity 2018 not applied.

Under H (columns 2-4): ocean models GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight (P+C formed per model, then averaged); measured transport error (GDP-replay OU fit per component, no added diffusivity); windage 0-5 % uniform; released 00:20 UTC 8 Mar.

Pleiades: rating-5 objects merged into 6 clusters at 3 km, equal cluster weights (rho4 = 0). COSMO-SkyMed: F1-F3, equal weights, dawn-20 Mar and dusk-21 Mar passes equally weighted; source, time and footprint unverfied.

Shading: density normalised to its maximum. Solid blue: 90 % HDR of the panel's PDF. Dashed grey: 90 % HDR of the flight posterior at the same stage. Circles: Pleiades rating-5 clusters; triangles: COSMO F1-F3. Grid 0.05 deg; 0.1-1.3 % of impact weight lies outside 85-103 E, 43-25 S and is not shown.

Every PDF in columns 2-4 is conditional on H (the objects are debris from 9M-MRO); no Bayes factor or P(H | data) is computed or implied.

Label: next-run (core (b), 4 strata mixed by P(family)); 00:19 option r1200/inflated; search Phase 2 + Bluefin-21; generated by prepare/branch_eof289.py + branch_figure.py (hypothesis/pleiades).