


P+C3 conditional under H after the base search (Phase 2 + Bluefin-21), by 00:19 option — next-run (core (b)) vs reference-289



Source: end of flight's next-run impacts on core (b) m0011 hand-offs (EoF 3c6319f, idle floor ON), 4 strata mixed with core's P(family) held fixed and not updated by terminal-stage evidence (free 0.6948, Davey dynamics + radar 0.1527, descent-climb 0.1376, routes 0.0149; core split-half 0.71-0.88 vs floor 0.896, unconverged); within each stratum 4 seeds pooled with equal weight, 3.2 M impacts per seed. Free stratum detail: impacts next-free (4 seeds pooled with equal weight; source run /Users/pete/claude-science/orgs/9db41e8b-db54-4736-82b9-d77e2a9ad222/workspaces/23b26fb9-3c70-4c8e-909b-ac78b132214a/runs/src-next-free, code 3c6319f); prior track 289.7 deg (sd 1.0); config chain (run.json): davey2016-inmarsat + no-exhaustion-prior + s1-factor + s2-temperature + s3-internal + s4-ceiling + s5-hard-reject + reference-snapshots + radar-full + free + server-free + hypotheses/end-of-flight/smoke/snapshot-m0011.toml + hypotheses/end-of-flight/smoke/terminal.toml + hypotheses/end-of-flight/full/reference-289.toml + hypotheses/end-of-flight/full/descent-idle-floor.toml + hypotheses/end-of-flight/full/seed-1.toml; tempered sampler (m1839, m1941, m2041, m2141, m2241, m0011, 16 stages); particles per mode [500000, 250000, 1250000, 250000, 1250000]; BFO bias (run.json bfo_bias): {"mean_hz":150.0,"sd_hz":25.0}.

Labels: core (b) split-half NOT converged; two-tank bookkeeping only; PROVISIONAL-OVERNIGHT; EoF sweep run by a stand-in; run by an architecture stand-in on the Pleiades module's behalf, module to review.

Recipe: Pleiades module's prepare/rerun_next.py at ed85311, unchanged, run per stratum (branch_eof289.py: weight = impact weight x exp(loglik:<option>), '+alive' adds end of flight's own constraint_log_factor, log-on cause 'other'); strata combined afterwards by the stand-in's combine_mixture.py with the module's own summarise(). Headline arm: rho4 = 0, equal cluster weights, GLORYS12 + GlobCurrent daily (+ ERA5) equal weight, P+C per model, measured spread.

Search: the searched-areas module's own per-impact seabed-search column (Phase 2 2014-17 + Bluefin-21, point target, rho = 0.05), from mh370 evaluate at ed85311 on these impacts.

Reference-289 markers: results/pleiades/branch-289/branch-by-0019-option.csv (plain options) and branch-289-alive/plain-vs-alive.csv (none, none+alive); eof-289-full, idle floor OFF. Options absent there have no marker. both/no-offset and both/startup-offset are parent-limited (52-265 effective parents per stratum): not estimates.

Every value is conditional on H (the objects are debris from 9M-MRO); no Bayes factor or P(H | data). PROVISIONAL-OVERNIGHT; unconverged upstream.