



Where the wreckage is on the seabed, for each use of the 00:19 satellite data

Flight families at the fixed weights of the flight model


(a) 00:19 Held Out




(b) 00:19 R600 BTO Only




(c) 00:19 R600 BTO + Raw BFO



(d) 00:19 Holland H1



(e) 00:19 Holland H2



- wreckage on the seabed, 50 % of probability
- wreckage on the seabed, 90 % of probability
- wreckage on the seabed, 99 % of probability
- - - impact on the water, 50/90/99 % of probability
- 6th arc, 00:11 UTC
- - - 7th arc, 00:19 UTC

Summary (STE): Grey areas show where the wreckage is on the seabed. Dashed orange lines show where the aircraft hit the water. The movement of the wreckage through the water adds less than 1 % to each grey area. The flight model is not yet stable, so these areas can change. There are not yet sufficient samples to show a result for Holland H1 and H2.

Technical: Settled-mass seabed density (real ocean: GLORYS12V1 column 75-115 E, 45-10 S, TEOS-10 density; GLORYS12V1 surface current; ERA5 wind; AusSeabed then GEBCO_2026), settling 5b595bf, breakup table PROVISIONAL, afloat pieces excluded. Impacts: end of flight next-run on core (b) (4 strata x 4 seeds, prior track 289.7 deg, EoF 43262c31), option_posteriors with +unpowered (airborne at 00:19:37, not powered at 01:15:56). Strata weighted by core's probability of each flight family only (fixed weights); P(family) for R600 BTO + Raw BFO = 0.695 / 0.153 / 0.138 / 0.015 (free / repro-radar / descent-climb / routes). Log-on cause: other (H1: fuel exhaustion). Systematic resample, one settling draw each: 200,000 (a), 40,000 (b-e). 0.02 deg grid, Gaussian 0.1 deg, HPD. Mixture ESS by Kish; below 1,000 not estimable (H1/H2: descent-model and sampler changes await Pete; architecture burst study). Labels: core (b) split-half NOT converged; two-tank bookkeeping only; internal-v1 one-engine flow 2x; dive class (b) and Boeing glide PROVISIONAL-OVERNIGHT; Mac, 2 threads.