


Where the wreckage is on the seabed, for each use of the 00:19 satellite data and each type of flight end

Rows: A1: fuel exhaustion, then no control; A2: fuel exhaustion, then controlled to the surface; B: deliberate descent before fuel exhaustion. Each panel is conditional on its row; the family share is the posterior probability of that row under the column's 00:19 data, not evidence for it.



Summary (STE): Grey areas show where the wreckage is on the seabed. Dashed orange lines show where the aircraft hit the water. Each row shows one type of flight end. The flight model is not yet stable, and the data hardly change how probable each type is, so do not read the shares as evidence. There are not yet sufficient samples for Holland H1 and H2.

Technical: Settled-mass seabed density, real ocean (GLORYS12V1 column 75-115 E, 45-10 S, TE05-10; GLORYS12V1 surface current; ERA5 wind; AusSeabed then GEBCO_2026), settling 9330d4c, breakup table PROVISIONAL, afloat pieces excluded. Impacts: end of flight on core (next-run-b), 4 strata x 4 seeds, prior track 289.7 deg, EoF ba26890, option_posteriors+unpowered; family codes (end of flight family_labels) A1 = [1.0], A2 = [2.0], B = [3.0] (family4_code, ruling 6 as amended: code 4 lost -> A2, undemonstrated recovery -> A1; code 6 outside B, share 10.8 %, 6.0 %, 7.6 %, 0.0 %, 0.7 % by column). Strata: P(stratum | option) (+alive) x share of the family. Systematic resample, one settling draw each: 60,000 (Held Out), 20,000 (others) per family. 0.02 deg grid, Gaussian 0.1 deg, HPD. Below 1,000 Kish ESS not estimable. Labels: core split-half NOT converged; two-tank bookkeeping only; dive class (b) PROVISIONAL; point mass cannot unload; Mac, 2 threads.