
If the imaged objects are from 9M-MRO: where the debris entered the sea — 00:19 R600 BTO + Raw BFO drift-error correlation 0.25, two ocean models (proposed headline) — core run C NOT CONVERGED




| | **ρ = 0.25 two models (drawn)** | ρ = 0.25 GLORYS12 only | ρ = 0 two models | ρ = 0.5 two models |
|---|---|---|---|---|
| 50 % region, km² | 7,457 | 5,817 | 7,511 | 7,530 |
| 90 % region, km² | 41,798 ± 247 | 30,998 ± 227 | 38,090 ± 229 | 44,951 ± 284 |
| mean position | 35.96 S 92.36 E | 36.01 S 92.38 E | 35.90 S 92.36 E | 36.00 S 92.36 E |
| outside past searches | 70.3 % | 74.3 % | 68.7 % | 71.2 % |
| share left by searches | 0.369 | 0.412 | 0.354 | 0.378 |
| tension ln S (p) | -0.30 (0.30) | +0.13 (0.43) | -0.48 (0.27) | -0.17 (0.33) |
| mean shift from PDF without H | 38 NM | 36 NM | 41 NM | 36 NM |
| PDF without H inside 90 % region | 16 % | 15 % | 15 % | 17 % |

STE100: This map shows where the aircraft possibly went into the sea, if the objects in the Pléiades satellite images are from 9M-MRO. The blue colour shows the probability with this assumption. The grey points show the probability without this assumption. The drawn result uses two ocean models with equal weight and a correlation of 0.25 between the drift errors of the two satellite data sets. The thin lines show the 90 % region for other settings. This is a proposed headline; the architecture session must accept it. The flight calculation (core run C) is not converged. Use these results with caution. Technical: core run C (4 strata x 8 seeds) split-half NOT CONVERGED (floor 0.924: free 0.918, Davey dynamics + radar 0.910, descent-climb 0.892, routes 0.850); end-of-flight physics PROVISIONAL (descents > 6,500 ft/min and unloading not reachable; no structural limit, G13). 00:19 R600 BTO + Raw BFO, existence constraint 'unpowered'; strata re-weighted by the 00:19 evidence (family-evidence-run-c.json, key '00:19 R600 BTO + Raw BFO +unpowered': next-c-free 0.606, next-c-repro-radar 0.218, next-c-descent-climb 0.154, next-c-routes 0.022). After Phase 2 + Bluefin-21 (official) + OI 2018 + OI 2025-26 SE band (grade C; NW band drawn only). Under H: 12 rating-5 Pléiades objects (6 clusters, 3 km, equal weight) x all four COSMO-SkyMed contacts, one debris field; bivariate-normal transport errors per component (prepare/rho_surface.py), ρ = 0.25 (ocean transport measured 0.07-0.41), ρ = 0 / 0.5 beside. GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight; GlobCurrent windage product-relative (drift audit F1: weights of c - dc, dc uniform 0.60-0.75 % U10 by 3-point quadrature, mass below 0 clamped to node 0; surfaces joint-surface-rho*-gcrel). PROPOSED headline (module, 11 Oct 18:19 UTC); in force until accepted: GLORYS12 only (ruling 2, 11 Oct 02:15 -0600), shown beside. ± = split-half σ of the 90 % region (rms over three 4 + 4 seed partitions, headline_splithalf.py). Conditional on H; no identity likelihood, no Bayes factor or P(H | data). Holland H1/H2 not estimable. Background: 6,000 points from the impact PDF without H after the same searches.