

Simplified English: The end-of-flight run now has extra samples of flights that end north of about 34 S. This chart shows the result of the seabed searches with and without these extra samples. The extra samples change the search result by a small amount (a). After the searches, more of the remaining probability is north of 33 S for most uses of the 00:19 data (b). In some northern bands the extra samples are still not sufficient when both 00:19 frequency values are used, mostly between 25 S and 20 S (c). Most of the remaining probability is either on searched seabed or more than 25 NM from the arc (d). A1 (cruise to fuel exhaustion, then no control) comes from two extra runs that sample only that type. The Ocean Infinity search areas come from community tracings and are a best estimate. Some parts of the flight-end physics are provisional.

Technical: Z = E[P(no find)]; P = rho + (1 - rho) prod_k (1 - q_k f_k c_k); base = ATSB Phase 2 (q 0.945) + Bluefin-21 (0.9) + OI 2018 (0.9, f 0.889) + OI 2025-26 outboard (0.9, f 0.7808), rho 0.05: Ocean Infinity coverage inferred (grade C), included as best estimate; planned next OI area not evidence. Source posterior: core run J (split-half accepted as a pass by project ruling, 16 Oct 2026), waypoint routes stratum only; end of flight rule B over all rows (n1 = 2,399,999 run J, n2 = 1,600,003 supplement, r_hat from core), reduced table impacts-ruleB.npz (2,408,038 rows, thinned with Horvitz-Thompson weights). A1 from end of flight's two conditional strata (no intervention: run J + first-half supplement pass; recovery attempted: run J paths only), end of flight's normalisation. Row mass: both log-on causes q = 1/3 with C2 silent, or 'other cause only' (C2 silent); D4/D5 Holland transient bias added. Uncontrolled-arm dynamics not calibrated; in-flight structural outcome PROVISIONAL. Grey: run J alone (all 153.6 M rows, A1 stratified, commit a8e819f) in (a); production alone on the reduced table in (b, c) and for the 'other cause only' rows. (b) seed bootstrap, 2,000 replicates, row weights fixed. (c) open = below 1,000 after the searches: both BFOs 35-30 S and 25-20 S, Holland bias 25-20 S, and 25-20 S under other cause only for both BFOs and Holland bias; paths = distinct hand-off rows per replicate (end of flight's distinct-state count is lower by up to 17 %). (d) impact point, arc distance from a 0.02 deg grid of end-of-flight values (p99 error 0.4 NM). Seabed: settling run-j-supplement (GLORYS12V1, ERA5, TEOS-10, AusSeabed/GEBCO_2026), field 'mean'. Drift, hydroacoustic and other evidence not used.