

Where the wreckage is on the seabed, for each route after 18:01 and each use of the 00:19 satellite data

Rows: one route each (named by its waypoints), in order of probability; each panel is conditional on its route


(a) Held Out · MEKAR-SAMAK-ISBIX




(b) R600 BTO Only · MEKAR-SAMAK-ISBIX




(c) R600 BTO + Raw BFO · MEKAR-SAMAK-ISBIX




(d) Holland H1 · MEKAR-SAMAK-ISBIX




(e) Holland H2 · MEKAR-SAMAK-ISBIX




(f) Held Out · NILAM-IGOGU-ISBIX




(g) R600 BTO Only · NILAM-IGOGU-ISBIX




(h) R600 BTO + Raw BFO · NILAM-IGOGU-ISBIX




(i) Holland H1 · NILAM-IGOGU-ISBIX




(j) Holland H2 · NILAM-IGOGU-ISBIX




(k) Held Out · NILAM-SAMAK-ISBIX




(l) R600 BTO Only · NILAM-SAMAK-ISBIX




(m) R600 BTO + Raw BFO · NILAM-SAMAK-ISBIX



(n) Holland H1 · NILAM-SAMAK-ISBIX



(o) Holland H2 · NILAM-SAMAK-ISBIX



(p) Held Out · NILAM-ANOKO-ISBIX

(q) R600 BTO Only · NILAM-ANOKO-ISBIX

(r) R600 BTO + Raw BFO · NILAM-ANOKO-ISBIX

(s) Holland H1 · NILAM-ANOKO-ISBIX

(t) Holland H2 · NILAM-ANOKO-ISBIX

wreckage on the seabed, 50 % of probability
wreckage on the seabed, 90 % of probability
wreckage on the seabed, 99 % of probability
impact on the water, 50/90/99 % of probability
6th arc, 00:11 UTC
7th arc, 00:19 UTC

Summary (STE): Grey areas show where the wreckage is on the seabed. Dashed orange lines show where the aircraft hit the water. The flight model is not yet stable, so these areas can change. In most Holland descents the aircraft would break up in the air before it hit the water; the model still puts the wreckage at one impact point.

Technical: Settled-mass seabed density (real ocean: GLORYS12V1 column 75-115 E, 45-10 S, TEOS-10 density; GLORYS12V1 surface current; ERA5 wind; AusSeabed then GEBCO_2026), settling 9330d4c, breakup table PROVISIONAL, afloat pieces excluded. Impacts: end of flight H1/H2 sweep on core run H (routes stratum) (1 strata x 8 seeds; EoF 7da43d01); row mass = core P(family) / seeds x hand-off weight x 00:19 likelihood (+unpowered; log-on lag density for H1) x (1 - P(in-flight breakup by 00:19:37)) [primary], as end of flight's holland_sweep_summary (no end-of-flight summary yet: In Z recorded, not checked); strata next-h-routes 1.000. Systematic resample over all rows, one settling draw each: 20,000. 0.02 deg grid, Gaussian 0.1 deg, HPD. Row ESS by Kish; below 1,000 not estimable. Labels: core run H = a data-selected 6-route subset (fly-by + leg tracking; 4 routes populated), not a fair family comparison with free flight; convergence not reported; piloted rapid descent to 40,000 ft/min with unloading to -1.0 g; INTERIM pitch-down; in-flight structural outcome PROVISIONAL (upper side: no transonic drag rise); breakup rows settled from the intact impact point (fragment field not modelled, G-bf); H1/H2 NOT estimable on run H (effective impacts 111 / 386); P(left) = 0.8 PROVISIONAL; Mac, 2 threads.