


Where the debris would have entered the sea if the imaged objects are from 9M-MRO — R1200, inflated BFO error



Source: impacts next-run (strata next-free, next-repro-radar, next-descent-climb, next-routes; provenance read from next-free) (4 seeds pooled with equal weight; source run src-next-free, code 3c6319f); prior track 289.7 deg (sd 1.0); config chain (run.json): davey2016-inmarsat + no-exhaustion-prior + s1-factor + s2-temperature + s3-internal + s4-ceiling + s5-hard-reject + reference-snapshots + radar-full + free + server-free + hypotheses/end-of-flight/smoke/snapshot-m0011.toml + hypotheses/end-of-flight/smoke/terminal.toml + hypotheses/end-of-flight/full/reference-289.toml + hypotheses/end-of-flight/full/descent-idle-floor.toml + hypotheses/end-of-flight/full/seed-1.toml; tempered sampler (m1839, m1941, m2041, m2141, m2241, m0011, 16 stages); particles per mode [500000, 250000, 1250000, 250000, 1250000]; BFO bias (run.json bfo_bias): {"mean_hz":150.0,"sd_hz":25.0}.

What was run: 00:19:37 log-on acknowledge (R1200) used; 00:19:29 message not used; BFO with an inflated, zero-mean error [end-of-flight option key: r1200/inflated]; log-on cause 'other'. Strata mixed by core's P(family), held fixed (next-free 0.6948, next-repro-radar 0.1527, next-descent-climb 0.1376, next-routes 0.0149); the search re-weights strata by their own evidence, the 00:19 data do not.

Labels: core (b): request-17 sampler fix IN, split-half NOT converged; two-tank bookkeeping only; internal-v1 fuel (its one-engine flow is 2x its tables); Inmarsat ephemeris; descent idle floor ON; EoF sweep and the Pléiades per-stratum branch run by architecture stand-ins and reviewed by the Pléiades module; PROVISIONAL-OVERNIGHT.

Search: the searched-areas module's own per-impact likelihood (point target, undetectable probability $\rho = 0.05$, shared between campaigns). Phase 2 (q 0.945) and Bluefin-21 (q 0.9) official; OI 2018 (q 0.9, coverage 0.889) and OI 2025-26 south-east band (q 0.9, coverage 0.7808) are inferred community tracings, grade C. The north-west band is drawn only and is never negative evidence.

Under H: the 12 rating-5 Pléiades objects merged into 6 clusters (3 km), equal weights; COSMO-SkyMed contacts equally weighted, dawn 20 Mar and dusk 21 Mar passes equally weighted (source, time and footprint unverified). 'One debris field' = both sets' likelihoods multiplied, per ocean model. Ocean models GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight; measured transport error (drifter replay, ~100-110 km per component at 15 d); windage 0-5 %; Pléiades and COSMO errors treated as independent ($\rho = 0$; 0.5-0.8 widened the 90 % area 28-38 % on reference-289).

Every PDF is conditional on H (the objects are debris from 9M-MRO); no Bayes factor or $P(H | \text{data})$. Shading: highest-density regions holding 50/90/99 % of each panel's PDF. Percentages are of the panel's PDF within 85-103 E, 43-25 S.