




Impact latitude with the Phase 2 and Bluefin searches as evidence






Where the probability goes (before: 30% on searched ground)




Detection probability q , and the inferred Ocean Infinity 2018 layer (labelled variant)






Davey eq. 11.2: probability of success, planning P_D 0.9



The best candidate areas, and what the search did to them



Seabed search evidence: numbers

IMPACTS FROM END-OF-FLIGHT MODULE END-OF-FLIGHT; 00:19 DATA OPTION NONE
Run eof-289-full: 4 replicates, 12,799,968 impacts; before: median -36.78, 95% -40.29 to -26.88; split-half 0.973; Kish ESS 12,358,799; on Phase 2 coverage 0.297

Mass removed at rho 0 (1 - evidence):
alone: bluefin-2014 0.0000
adding, in order: bluefin-2014 0.0000
alone: phase2-2014-2017 0.2805
adding, in order: phase2-2014-2017 0.2805
alone: ocean-infinity-2018 0.0400
adding, in order: ocean-infinity-2018 0.3202

scenario	Z	median	q025	q975	N of 33S	S of 39.5	on P2	ESS	split	vs before
run.toml	0.7335	-36.58	-40.51	-26.29	0.138	0.094	0.044	9,477,480	0.968	0.879
rho 0	0.7195	-36.56	-40.53	-26.26	0.140	0.096	0.026	9,150,141	0.968	0.870
rho 0.02	0.7251	-36.57	-40.52	-26.27	0.139	0.095	0.033	9,282,628	0.968	0.873
rho 0.1	0.7476	-36.60	-40.50	-26.33	0.135	0.092	0.062	9,791,006	0.968	0.887
rho 0.2	0.7756	-36.63	-40.47	-26.39	0.130	0.089	0.096	10,370,668	0.969	0.903
rho 0.3	0.8037	-36.66	-40.45	-26.45	0.126	0.086	0.127	10,879,410	0.970	0.918
rho 0.5	0.8598	-36.71	-40.40	-26.58	0.118	0.080	0.183	11,663,865	0.971	0.946
Phase 2 q 0.90	0.7462	-36.60	-40.50	-26.32	0.135	0.093	0.060	9,761,782	0.968	0.887
Phase 2 q 0.98	0.7237	-36.57	-40.52	-26.27	0.140	0.095	0.031	9,248,475	0.968	0.873
Phase 2 split, shared misses	0.7318	-36.58	-40.52	-26.29	0.138	0.094	0.042	9,436,515	0.968	0.878
Phase 2 split, independent misses	0.7301	-36.58	-40.52	-26.28	0.138	0.095	0.040	9,394,909	0.968	0.877
+ OI 2025-26 inferred, outboard band only	0.7298	-36.60	-40.52	-26.28	0.138	0.095	0.044	9,448,274	0.968	0.879
+ OI 2018 inferred, coverage 0.889	0.6958	-36.76	-40.55	-26.35	0.120	0.099	0.046	9,121,248	0.968	0.898
+ OI 2018 inferred, coverage 0.952	0.6931	-36.77	-40.56	-26.35	0.118	0.100	0.046	9,067,284	0.968	0.899

Residual PDF view. Source posterior: eof-289-full, 00:19 option none; search evidence applied with run.toml. Davey eq. 11.2 at a planning P_D of 0.9 on 0.5 deg blocks:

rank	candidate	area	residual mass	P(find)	km2	same block, search disabled
1	-35.50	91.00 E 0.50 deg	0.0235	0.0212	2524	0.0172
2	-35.00	91.50 E 0.50 deg	0.0189	0.0170	2540	0.0139
3	-36.50	89.50 E 0.50 deg	0.0178	0.0160	2493	0.0141
4	-37.00	89.00 E 0.50 deg	0.0155	0.0140	2477	0.0166
5	-36.00	90.50 E 0.50 deg	0.0153	0.0138	2509	0.0152
6	-36.00	90.00 E 0.50 deg	0.0144	0.0130	2509	0.0106
7	-37.00	88.50 E 0.50 deg	0.0141	0.0127	2477	0.0105
8	-39.00	89.00 E 0.50 deg	0.0136	0.0122	2411	0.0100

P(find) 25% needs the best 21 blocks, 51,977 km2
P(find) 50% needs the best 62 blocks, 152,587 km2
P(find) 75% needs the best 231 blocks, 580,334 km2