Where the wreckage is on the seabed, by type of flight end (waypoint routes, with the extra northern flights)

Rows: A1 cruise to fuel exhaustion, then no control; A2 cruise to fuel exhaustion, then controlled descent; B planned descent with ditching approach begun before fuel exhaustion

(a) Held Out · A1
90 % area (thousand km²): seabed 36; impacts 36
10,000 draws, 10,000 impacts; ESS 41,685
share of the treatment 17.0 %

(b) R600 BTO Only · A1
90 % area (thousand km²): seabed 21; impacts 21
10,000 draws, 10,000 impacts; ESS 38,454
share of the treatment 19.6 %

(c) R600 BTO + Raw BFO · A1
90 % area (thousand km²): seabed 20; impacts 20
10,000 draws, 9,960 impacts; ESS 33,250
share of the treatment 23.6 %

(d) both BFOs as recorded · A1
Not yet estimable
row ESS 408 (< 1,000)
90 % area (thousand km²): seabed 18; impacts 18
10,000 draws, 1,539 impacts; ESS 408
share of the treatment 9.3 %

(e) both BFOs, Holland transient bias added · A1
90 % area (thousand km²): seabed 27; impacts 27
10,000 draws, 7,358 impacts; ESS 2,913
share of the treatment 24.6 %

(f) R1200 BFO, Holland transient bias added · A1
90 % area (thousand km²): seabed 18; impacts 18
10,000 draws, 2,840 impacts; ESS 1,455
share of the treatment 5.8 %

(g) Held Out · A2
90 % area (thousand km²): seabed 166; impacts 166
10,000 draws, 10,000 impacts; ESS 77,395
share of the treatment 34.1 %

(h) R600 BTO Only · A2
90 % area (thousand km²): seabed 125; impacts 125
10,000 draws, 10,000 impacts; ESS 68,349
share of the treatment 34.5 %

(i) R600 BTO + Raw BFO · A2
90 % area (thousand km²): seabed 130; impacts 130
10,000 draws, 9,630 impacts; ESS 24,099
share of the treatment 23.3 %

(j) both BFOs as recorded · A2
90 % area (thousand km²): seabed 59; impacts 59
10,000 draws, 7,779 impacts; ESS 4,451
share of the treatment 40.6 %

(k) both BFOs, Holland transient bias added · A2
90 % area (thousand km²): seabed 59; impacts 59
10,000 draws, 9,875 impacts; ESS 30,460
share of the treatment 34.8 %

(l) R1200 BFO, Holland transient bias added · A2
90 % area (thousand km²): seabed 57; impacts 57
10,000 draws, 9,318 impacts; ESS 14,195
share of the treatment 41.7 %

(m) Held Out · B
90 % area (thousand km²): seabed 188; impacts 187
10,000 draws, 10,000 impacts; ESS 114,317
share of the treatment 48.8 %

(n) R600 BTO Only · B
90 % area (thousand km²): seabed 134; impacts 134
10,000 draws, 10,000 impacts; ESS 93,685
share of the treatment 45.8 %

(o) R600 BTO + Raw BFO · B
90 % area (thousand km²): seabed 116; impacts 115
10,000 draws, 9,722 impacts; ESS 35,611
share of the treatment 53.0 %

(p) both BFOs as recorded · B
90 % area (thousand km²): seabed 69; impacts 69
10,000 draws, 6,863 impacts; ESS 1,968
share of the treatment 50.0 %

(q) both BFOs, Holland transient bias added · B
90 % area (thousand km²): seabed 66; impacts 66
10,000 draws, 9,626 impacts; ESS 16,069
share of the treatment 40.4 %

(r) R1200 BFO, Holland transient bias added · B
90 % area (thousand km²): seabed 64; impacts 64
10,000 draws, 8,000 impacts; ESS 7,173
share of the treatment 52.4 %

latitude: 20°S, 25°S, 30°S, 35°S, 40°S
longitude (°E): 85, 90, 95, 100, 105

wreckage on the seabed, 50 % of probability
wreckage on the seabed, 90 % of probability
wreckage on the seabed, 99 % of probability
impact on the water, 50/90/99 % of probability
6th arc, 00:11 UTC
7th arc, 00:19 UTC

Summary (STE): Grey areas show where the wreckage is on the seabed. Dashed orange lines show where the aircraft hit the water. Each map uses one way to read the 00:19 satellite data. The flight model now has more flights that end in the north; a known weight keeps the probabilities correct. Some parts of the flight model are provisional, so these areas can change. When the aircraft breaks up in the air, the model still puts the wreckage at one impact point. Search results and debris drift are not used here.

Technical: Settled-mass seabed density (real ocean: GLORYS12V1 column 75-115 E, 45-10 S, TEOS-10; GLORYS12V1 surface current; ERA5 wind; AusSeabed then GEBCO_2026), settling 9330d4c, breakup table PROVISIONAL, afloat pieces excluded. Impacts: end of flight's rule-B reduced table (impacts-ruleB.npz, 2,408,038 rows): core run J waypoints production (24 seeds) + northern supplement (16 seeds, all rows) + extra descents (D2-D5), combined by rule B with core's r_hat, one Horvitz-Thompson thinning; binary mh370-eof-rj1. Waypoints stratum only (conditional on it). Row weight = end of flight's stored lnw:comb (hand-off weight x survival x L(00:19 treatment) x L(cause), both causes q = 1/3 with C2 silent, W = 00:10:58-01:15:56; 'other log-on cause only' = C2-silent arm). D3 = R600 BTO + both BFOs as recorded; D4 = both BFOs, Holland transient bias added; D5 = R600 BFO as recorded + R1200 BFO, Holland transient bias added U[17, 130] Hz; BFO error 4.3177 Hz. Systematic resample, one settling draw each: 10,000. 0.02 deg grid, Gaussian 0.1 deg, HPD. Row ESS by Kish; below 1,000 not estimable. Held Out is D0' (impact after 00:19:37.443 imposed). Prior track: core run J waypoint routes. Labels: core run J split-half accepted as a pass by project ruling, 16 Oct 2026; uncontrolled-arm dynamics not calibrated (Boeing leave-one-out gate failed); piloted-turn model and INTERIM pitch-down PROVISIONAL; in-flight structural outcome PROVISIONAL; breakup rows settled from the intact impact point (G-bf); P(left) = 0.8 PROVISIONAL; Mac.