
Seabed wreckage latitude: the extra northern flights fill in the north of the 7th arc and do not change the southern peak



(i) The lines show where the wreckage is on the seabed, by latitude, for each way to read the 00:19 satellite data (all types of flight end; both log-on causes, fuel exhaustion 3 : 1 other). The dashed grey line uses the main flights only. The blue line also uses extra flights that start in the north, with a known weight that keeps the probabilities correct. North of 33° S (thin vertical line) the blue line comes from a separate sample of northern impacts. The light grey line is the published Davey et al. (2016) curve (drawn where it is above 0.001 per degree). The vertical scale is logarithmic. Some parts of the flight model are provisional. Search results and debris drift are not used.

(ii) Settled-mass-weighted seabed latitude, 0.1° bins, afloat pieces excluded; settling 9330d4c (GLORYS12V1, TEOS-10, ERA5, AusSeabed/GEBCO_2026; breakup table PROVISIONAL; breakup rows from the intact impact point, G-bf). Main flights: settling run J waypoints (24 seeds, 40,000 outcomes). With the extra flights: end of flight's rule-B reduced table (run J production + all 16 supplement seeds + extra descents for D2-D5, core r_hat), 40,000 outcomes, and north of 33° S a conditional product of 10,000 outcomes scaled by its share. Prior track: core run J waypoint routes (stratum conditional, core share 0.9982). Primary cell: q = 1/3, C2 silent; Held Out is D0'. BFO error 4.3177 Hz; the main-flights D5 curve uses the 18 Oct table (D5 ln Z differs by 0.018). Uncontrolled-arm dynamics not calibrated. Core run J split-half accepted as a pass by project ruling, 16 Oct 2026. Davey et al. (2016), Fig. 10.3 recovered from the PDF vector paths (a flight posterior, shown for reference).