
Where the wreckage is on the seabed (core run I, not converged)
Dark: 50 % region of the wreckage. Light: 90 % region. Dashed line: 90 % region of the impact point.



(i) The wreckage moves very little as it sinks, so the seabed regions are almost the same as the impact regions. The right column is correct only if the Pléiades objects are debris. Breakup in the air is not modelled for the debris field. Do not use these maps for a search.
(ii) Technical: Core run I: 4 core families x 8 seeds x 100,000 hand-off rows, equal family prior; NOT CONVERGED (split-half waypoints 0.916, floor 0.924; free flight 0.830, descent-climb 0.794, Davey reproduction 0.891). End of flight run I (binary mh370-eof-ri1, 102.4 M impacts): piloted turns and interim pitch-down provisional; no structural limit (G13). 00:19: competing log-on causes q = 1/3, W = 00:10:58-01:15:56, C2 silent, survival factor applied; no treatment uses the R1200 BTO. Ocean: GLORYS12 and GlobCurrent (product-relative windage), equal prior, mixed per row (Bayesian model averaging). Drift h50; unscored rows carried at the mean ratio (<= 0.11 % of weight); Western Australia non-recovery not modelled (G26). Hydroacoustics predictive only, not in these products. Searches: ATSB Phase 2 + Bluefin-21, rho 0.05, point target; Ocean Infinity excluded. Pléiades: rho 0.25, rating 5, six clusters plus four COSMO-SkyMed contacts, conditional on the objects being debris. Strata mixed by core share x 00:19 evidence (ruling C), conditional on equal family prior. Composer pass 2, architecture stand-in. Seabed: settling run I samples (settling 9330d4c; 40,000 outcomes per stratum and treatment, from the flight + end-of-flight weights), re-weighted per outcome by the composed module factors (importance weights; outcome ESS 1,300-1,900 in the right column); settled mass only, afloat 18 %; breakup rows settled from the intact impact point (G-bf). Strata by settling's core share x Z(cell).