


What the chart shows. Each square is a possible impact point; the points are 30 NM apart. The colour shows how well the drift of debris from that point explains the nine identified debris finds. Panels a and b use the same scale. Panel c shows where the two ocean models agree. Panel d shows which model gives the higher value. The GlobCurrent run now uses a smaller wind effect, because its surface current already includes some wind drift. Compare with the earlier chart, where the two models did not agree. The results are interim and are not evidence.

Technical. Runs debris-drift-production-glorys12 (merged; binary d24060aa8006d3ce) and debris-drift-runA-globcurrent (merged, 8 chunks; binary e6bb17fe32f22b55; 710 nodes, the 367 planned nodes shown), Darwin arm64; 10^5 particles/node (3 windage classes × 8,334 × 4 ocean-error realisations, seeds 1-4); GLORYS12V1 0.494 m and GlobCurrent my-p1d 0 m, each + ERA5 10 m wind; GSHHG coastline; recovery bandwidth 50 km; node extent reference-289 (prior track 289.7°). Log-likelihood relative to joint maximum (-125.19). c: −|Δ log-likelihood|; outlined where |Δ| ≤ 2σ of combined split-half Monte Carlo noise (σ 1.42 GLORYS12, 1.09 GlobCurrent). Node correlation 0.39. GlobCurrent windage product-relative: c_wind offset drawn per particle U(−0.75%, −0.60%) of U10 (audit F1); before the correction, correlation −0.08 and 293 of 367 within 2σ. 00:19 option not applicable (per-node surface).