
If the imaged objects are from 9M-MRO: where the debris entered the sea — 00:19 R600 BTO + both BFOs (Holland H2 data)
core run I, drift-error correlation 0.25, two ocean models (proposed headline) — NOT CONVERGED — NOT ESTIMABLE (too few samples under H)




| | **ρ = 0.25 two models (drawn)** | ρ = 0.25 GLORYS12 only | ρ = 0 two models | ρ = 0.5 two models |
|---|---|---|---|---|
| 50 % region, km² | 1,130 | 877 | 1,054 | 1,179 |
| 90 % region, km² | 7,644 | 6,635 | 6,739 | 8,698 |
| mean position | 35.68 S<br>92.56 E | 35.74 S<br>92.55 E | 35.65 S<br>92.57 E | 35.70 S<br>92.54 E |
| outside official searches | 79.4 % | 81.7 % | 78.9 % | 79.4 % |
| share left by searches | 0.417 | 0.465 | 0.410 | 0.417 |
| tension ln S (p) | -0.03 (0.22) | +0.32 (n/e) | -0.12 (0.24) | +0.04 (0.20) |
| mean shift from PDF without H | 81 NM | 79 NM | 83 NM | 80 NM |
| PDF without H inside 90 % region | 21 % | 20 % | 19 % | 23 % |

- probability density if the objects are debris (relative)
- 50 % region (drawn setting)
- 90 % region (drawn setting)
- 90 % region, drift-error correlation 0.25, GLORYS12 only
- 90 % region, drift-error correlation 0, two ocean models
- 90 % region, drift-error correlation 0.5, two ocean models
- impact PDF without H (sampled points)
- ATSB Phase 2 2014-17 (official)
- Ocean Infinity 2018 (grade C) — drawn only, not applied
- OI 2025-26 south-east band (grade C) — drawn only, not applied
- OI north-west band (not searched; grade C)
- 7th arc, FL400
- Pléiades rating-5 objects
- COSMO-SkyMed radar contacts (all four)

STE100: This map shows where the aircraft possibly went into the sea, if the objects in the Pléiades satellite images are from 9M-MRO. The blue colour shows the probability with this assumption. The grey points show the probability without this assumption. The drawn result uses two ocean models with equal weight and a correlation of 0.25 between the drift errors of the two satellite data sets; the thin lines show the 90 % region for other settings. Only the official searches (Phase 2 and Bluefin-21) are applied. The flight calculation (core run I) is not converged. Use these results with caution. Technical: core run I (4 strata x 8 seeds; waypoints carries core P 0.9982) NOT CONVERGED; end-of-flight physics PROVISIONAL (piloted-turn model, INTERIM pitch-down, push-over and Boeing sequence; end of flight's run I labels). 00:19 R600 BTO + both BFOs (Holland H2 data); log-on cause: competing-causes total, q = 1/3, C2 silent (primary); survival factor applied; end of flight's own row weights (option_table_0019), strata by core P x own evidence. Search: searched areas' per-impact ln P(no find), rho 0.05, point target, Phase 2 (q 0.945) + Bluefin-21 (q 0.9); OI 2018 and 2025-26 not applied (grade C; drawn only). Under H: 12 rating-5 Pléiades objects (6 clusters, 3 km, equal weight) x all four COSMO-SkyMed contacts, one debris field; bivariate-normal transport errors, ρ = 0.25 (measured 0.07-0.41), ρ = 0 / 0.5 beside; GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight, GlobCurrent windage product-relative (drift audit F1). PROPOSED headline; GLORYS12 only (ruling 2) beside. Maps from per-impact columns on the wide 0.05 deg grid (prepare/runi_closeups.py). Conditional on H; no identity likelihood, no Bayes factor or P(H | data). Coverage: effective impacts after the official searches 115 under H (656 without H); A1 under H 25. NOT ESTIMABLE: below the 1,000 gate. The speckled regions, the small areas and the tension are sampling artefacts of too few effective impacts, not a measured result. Background: 6,000 points from the impact PDF without H after the same searches.