The searches remove more of the probability on run I, because run I puts more of it where the sonars looked



WHAT THIS SHOWS. Left: how much of the probability for where the aircraft hit the water lies on sea floor the ATSB had already searched, for each of the five ways of reading the two radio transmissions at 00:19 UTC. Grey is the previous set of simulated flights; dark blue is the newest set, which uses a new model of how the aircraft turned. Right: what the searches, which found nothing, then do to that probability. Because the newest flights place more of it where the sonars looked, the searches now take away about half of it instead of about a third. The four markers are the four families of flight path; the filled one carries almost all of the weight. The flights themselves are not yet stable, so treat all of this as provisional.

TECHNICAL. Source: end of flight run I, 4 strata x 8 seeds, 102,399,904 impacts, binary mh370-eof-ri1 (aa79d44e); against run D-he, 4 strata x 4 seeds, 51,200,064 impacts. Primary arm (survival factor 1 - P(in-flight breakup before 00:19:37) applied to the posterior weights), constraint +alive. NO MIXTURE IS SHOWN FOR RUN I: core has published run I P(family) only between free flight and waypoints (ln BF +6.78, so waypoints take 0.999 of those two), not the four-family share, so each stratum is plotted separately and none is weighted. Run D-he uses end of flight's holland-sweep-summary.json weights 0.594 / 0.235 / 0.153 / 0.019. One mh370 evaluate pass per seed over eight coverage layers; each scenario is exact algebra on the raw coverage columns, checked against evaluate's own no_find_probability on every seed (max |delta| < 1e-12). Likelihood P(no find | y) = rho + (1 - rho) prod_k [1 - c_k(y) g_k q_k], point target, shared miss dependence, raster 0.01 deg bilinear. ATSB Phase 2 union 120,486.5 km2 at q 0.945; Bluefin-21/Artemis 771.4 km2 at q 0.900 (removes 0.0000 at every rho); Ocean Infinity layers excluded. rho = 0.05, NO PUBLISHED VALUE: sweeping it 0 to 0.5 moves run I waypoints Z by 0.26 (Held Out) to 0.41 (Holland H1). LABELS CORE RUN I NOT CONVERGED (core waypoints split-half 0.916); this module's Z split-half gap is 0.040-0.044 on run I against 0.003-0.008 on run C, so the third digit is noise. End of flight physics PROVISIONAL: piloted turns (Davey turns with each particle's own tau_h, then a 25-minute terminal window of random turns, sweep and ditching circuit), INTERIM parametric pitch-down, aimed pitch-down for A1, push-over to 40,000 ft/min at VMO 330 KCAS, C2-silent proposal component, residual roll P(left) = 0.8, trim at loss and Boeing sequence applied. Wreckage field NOT applied on run I (settling has posted no run I samples): point target only. COVERAGE GAP: family A1 within Holland H1/H2 is below the 1,000 gate under end of flight's D3 and D5 cells (351 / 252 effective impacts).