


The best-placed archived infrared images of the most probable aircraft area (two best overall, best from a second satellite), 7 March 2014

FY-2D full disc, scan start 19:31:00
10.8 µm brightness temperature; aircraft reached 19:44 UTC




FY-2D full disc, scan start 19:31:00
10.8 minus 12.0 µm difference; aircraft reached 19:44 UTC







FY-2D full disc, scan start 21:01:00
10.8 µm brightness temperature; aircraft reached 21:17 UTC




FY-2D full disc, scan start 21:01:00
10.8 minus 12.0 µm difference; aircraft reached 21:17 UTC







MTSAT-2 full disc, scan start 18:31:15
10.8 µm brightness temperature; aircraft reached 18:42 UTC



MTSAT-2 full disc, scan start 18:31:15
10.8 minus 12.0 µm difference; aircraft reached 18:42 UTC





Plain note: These are the two archived images that saw the most probable aircraft area most clearly (clear sky, small pixels), and the best image from a second satellite. Weather analysis shows the air along these routes was too dry for a lasting condensation trail, so none is expected; the images are shown as a check, not as a search.

Technical note: ranking = weighted fraction of posterior apparent positions over 10.8 um > 285 K divided by the effective IR pixel size at the median position (geo_top_pass_ranking.csv). Shown: the two top-ranked scans, FY-2D 19:31 (rank 1; VZA 9.0 deg, 5.0 km) and FY-2D 21:01 (rank 2; 18.6 deg, 5.2 km), plus the best-ranked MTSAT-2 scan, 18:31 (rank 5; 58.4 deg, 5.9 km), so that a second satellite is represented; FY-2D 21:31 (rank 3) and 20:31 (rank 4) are in the supplementary FY-2D figure. Posterior: core run J BTO+BFO, 148,000 routes, family weights waypoints 0.9982, free 0.0013, repro-radar 0.0003, descent-climb 0.0002, FL350, position at the modelled scan time; highest-density regions from a 0.1 deg weighted histogram. ERA5 along these routes: no ice supersaturation at FL300-FL400 (posterior probability of a persistent contrail < 1e-6; contrail-notes.md, artifact df5975a4). Sources as in the sensor table.