




Source: end of flight's reference-289 impacts (run eof-289-full; core reference-289, prior track 289.7 deg at 18:01:49 UTC), 4 seeds pooled with equal weight, 12,799,968 impacts. 00:19 data option 'r600/no-offset' (project priority 2, R600 only, raw): the 00:19:29.416 R600 log-on request's BTO and BFO score the impacts with no offset manipulation; R1200 held out; log-on cause 'other' (no fuel-exhaustion log-on density). Weight = impact weight x exp(loglik:r600/no-offset).

End-of-flight physics PROVISIONAL: dive class (b) (being rebuilt against Boeing's simulator runs); Boeing-calibrated glide (Pete's reference).

Search (row 2): the searched-areas module's own per-impact likelihood, base case Phase 2 2014-17 + Bluefin-21, point target, undetectable probability $\rho = 0.05$; Ocean Infinity 2018 not applied.

Under H (columns 2-4): ocean models GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight (P+C formed per model, then averaged); measured transport error (GDP-replay OU fit per component, no added diffusivity); windage 0-5 % uniform; released 00:20 UTC 8 Mar.

Pleiades: rating-5 objects merged into 6 clusters at 3 km, equal cluster weights ($\rho_4 = 0$). COSMO-SkyMed: F1-F3, equal weights, dawn-20 Mar and dusk-21 Mar passes equally weighted; source, time and footprint unverified.

Shading: density normalised to its maximum. Solid blue: 90 % HDR of the panel's PDF. Dashed grey: 90 % HDR of the flight posterior at the same stage. Circles: Pleiades rating-5 clusters; triangles: COSMO F1-F3. Grid 0.05 deg; 1.5-2.0 % of impact weight lies outside 85-103 E, 43-25 S and is not shown.

Every PDF in columns 2-4 is conditional on H (the objects are debris from 9M-MRO); no Bayes factor or $P(H \mid \text{data})$ is computed or implied.

Label: 289.7 prior (reference-289); EoF impacts, 00:19 option r600/no-offset (R600 only, raw); generated by prepare/branch_eof289.py + branch_figure.py (hypothesis/pleiades).