





c

Tension by descent kernel

● 289.7° prior ● 295.66° prior



Panels a-b: core reference-289 (prior track 289.7 deg, 4 seeds x 7M particles), per-particle positions at 00:19:37 UTC, convolved with the PROVISIONAL eof-2f descent kernel. No 00:19 burst scoring of impacts and no seabed-search evidence.

Panel c: $\ln S$ (Handley & Lemos 2019), pooled (dot) and seed range (bar), per descent kernel: disks; eof-2f; end of flight's displacement histograms (Boeing glide, +/-160 NM, SMOKE scale, seed 1, N = 16, made on the 295.66 hand-off; 'none' = 00:19 messages held out, r600/r1200 = 00:19:29 / 00:19:37 message scored; 'other' / 'fuel-exhaustion' = log-on cause; dive class (b) on/off, PROVISIONAL).

Under H: ocean models GLORYS12 + ERA5 and GlobCurrent daily + ERA5 at equal weight; measured transport error (GDP-replay OU fit per component, no added diffusivity); windage 0-5 % uniform; released 00:20 UTC 8 Mar. Pleiades rating-5 objects in 6 clusters at 3 km, equal cluster weights ($\rho_4 = 0$). The conditional is $p(\text{impact} \mid \text{Pleiades}, H)$; no Bayes factor or $P(H \mid \text{data})$ is computed or implied.