




Prior: end of flight's reference-289 impacts (run eof-289-full; core reference-289, prior track 289.7 deg sd 1.0 at 18:01:49 UTC; base config davey2016 + no-exhaustion-prior + reference-snapshots + early-families/overnight/reference-289 + end-of-flight full/reference-289); 4 seeds x 100,000 parents x 8 children x 4 descents = 12,799,968 impacts; 7 Hz BFO sigma baseline. 00:19 option 'none' (both 00:19 messages held out), log-on cause 'other'. End-of-flight dive class (b) and Boeing-calibrated glide: PROVISIONAL.

Search: the searched-areas module's per-impact likelihood, point target, rho = 0.05. Phase 2 (q 0.945) and Bluefin-21 (q 0.9) official; OI 2018 (q 0.9, coverage fraction 0.889) and OI 2025-26 south-east band (q 0.9, coverage fraction 0.7808) are inferred community tracings (MH370-CAPTION, grade C), used as Pete ruled with this footnote. The north-west band is drawn only: it is not searched and is never negative evidence.

Under H: Pléiades rating-5 clusters (3 km), equal weights; COSMO pass dawn-20 / dusk-21 Mar equally weighted; P+C = both sets debris (product), formed per ocean model; GLORYS12 + ERA5 and GlobCurrent daily + ERA5 at equal weight; measured transport error (GDP-replay OU fit), windage 0-5 %. Conditional on H; no Bayes factor or P(H | data). Shading normalised per panel. Percentages are of the panel's PDF within the 0.05 deg grid.