Fuel exhaustion then no control, under the satellite-object hypothesis: with all supplement paths, the cells with the Holland transient bias added pass 1,000; the other four cells stay below it (409-702)



Footnote (simple): The data is conditional on the hypothesis that one or more objects seen by satellite on 20-23 March 2014 came from the aircraft. We do not use the shape of the objects. The end of flight physics is provisional; the no-control descent model is not calibrated. No seabed search is applied. The family is: cruise to fuel exhaustion, then no control. 20 October adds all the northern supplement paths and a second sample of the descents with an attempted recovery. The gate is 1,000 effective impacts.
Footnote (technical): Core run J (split-half accepted as a pass by project ruling, 16 Oct 2026), waypoints stratum; rule B over 2,399,999 run J and 1,600,003 supplement hand-off rows (r_hat from core), extra descents 64:256 (D2-D5). A1 = branch 1 (no intervention; 18 Oct pass + 19 Oct supplement pass, rule B) + branch 2 A1 rows (recovery attempted; 20 Oct pass, run J paths), end of flight's stratified estimator with L_H on every row; Kish ESS. H: Pleiades rating 5 (6 clusters) + COSMO-SkyMed F1-F4, pass-marginal; rho 0.25; GLORYS12+ERA5 and GlobCurrent+ERA5 (product-relative windage), equal prior. 19 Oct D5 used the 7 Hz BFO table; 20 Oct D5 uses 4.3177 Hz (end of flight). No identity likelihood; no Bayes factor or P(H). Before any seabed search.