
Wreckage on the seabed if the Pléiades objects were aircraft debris

The impacts that go to the seabed calculation are drawn from the weights of each combination of evidence



Summary (STE): Grey areas show where the wreckage is on the seabed. Dashed orange lines show where the aircraft hit the water. Each panel uses the weights of one combination of evidence: debris = where floating debris was found; sound = underwater sound records; Pléiades = if the objects seen in the Pléiades satellite images were aircraft debris; searches = the areas searched without a find. The flight model and the other evidence are not yet stable, so these areas can change.

Technical: Settled-mass seabed density (real ocean: GLORYS12V1 column 75-115 E, 45-10 S, TEOS-10 density; GLORYS12V1 surface current; ERA5 wind; AusSeabed then GEBCO_2026), settling 9330d4c, breakup table PROVISIONAL, afloat pieces excluded. Impacts: end of flight on core run C (4 strata x 8 seeds, EoF 87f6343); per-row weights from the composer (product per panel; constraint unpowered; composer pass 1 on run C (architecture stand-in): H = flight model x end of flight x debris drift (GLORYS12V1) x hydroacoustics stand-in L_hyd x Pléiades (rating 5, equal weights, GLORYS12V1), CONDITIONAL on the Pléiades objects being aircraft debris; the second form also x searched areas (rho 0.05); all rows, no speed cap; GlobCurrent excluded (drift audit F1); PIPELINE TEST). Systematic resample over all rows, one settling draw each: 40,000, 40,000, 40,000, 40,000. 0.02 deg grid, Gaussian 0.1 deg, HPD. Row ESS by Kish; below 1,000 not estimable. Labels: core run C split-half NOT converged (min 0.82-0.86 against floor 0.924); rapid descents above 6,500 ft/min and unloading not reachable; residual roll P(left) = 0.8 PROVISIONAL; trim at loss and two-tank takeover NOT applied; Mac, 2 threads.