



MH370 posterior evolution and seabed-search context

00:11 medium-BFO posterior → corrected-R600 00:19 control → primary conditional v0.1 impact family


A Posterior evolution and southern search context



B Davey 00:19 latitude PDF



C Full searched-area locator



D Search-area evidence

- Phase 2 official display geometry
- OI 2018 approximate outline (C)
- Renewed outboard estimate (C)
- Renewed inboard estimate (C)
- AIS surface-vessel proxy (C)
- Official seventh arc
- Bluefin official display polygons

Mapped / reported area context

Phase 2 catalog: Deep Tow 130,961; DHJ 46,884;
GO Phoenix 82,606 km² (overlap; not additive)
Bluefin 771 km² display / 860 km² reported
OI 2018 >112,000 km² reported;
exact swaths unplotted
OI 2025-26 7,571 km² surveyed;
7,429 km² arithmetic remainder only

CONTEXT ONLY — no binary searched/not-searched mask.
No calibrated target-specific probability of detection.
Grade-C OI outlines and AIS tracks are educated context,
not AUV sonar swaths.

Posterior contours are separate 50/95/99% display HPDs on a 4 km local metric raster with a fixed 7.5 NM Gaussian kernel; numerical weights are unchanged. The states are sequential and dependent. Search geometry is contextual only; exact OI/Bluefin swaths, holidays, quality and calibrated PoD remain unresolved.