

MH370 flaperon drift diagnostic - CMEMS GLORYS12

Uncalibrated family-specific diagnostic; currents=cmems_glorys12_reanalysis | isotope=enabled as a neutral-or-penalizing sc

Estimate at a glance

CURRENT FAMILY
cmems_glorys12_reanalysis

PEAK POSITION
-16.195 deg , 106.100 deg

MEAN IMPORTANCE-WEIGHTED FIELD-TERMINATION FRACTION
1.4%

ARRIVAL SAMPLER
AdaptiveSmc


PEAK SOURCE CELL
arc-126

WEIGHTED TRAJECTORIES
2048 ? 77 cells

MEAN PROPOSAL TERMINATION FRACTION
0.6%

PEAK CORRECTION ESS
973.3


dark outline: 50% HDR
blue outline: 90% HDR



Probability is conditional on the named motion, SATCOM, and evidence configuration.

Posterior geography

Highest-density regions are computed from the weighted particle population.




Interpretation

The map is a probability density over the modelled impact state. It is not a searched-area elimination map, and structural model alternatives must be compared separately.

Numerical behaviour and conditions

Adaptive likelihood tempering, particle diversity, and declared evidence conditions.



Evidence conditions

- Current family selected explicitly: cmems_glorys12_reanalysis
- Barnacle isotope: enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival
- direct windage disabled
- Stokes drift omitted
- explicit land fraction and calibrated beaching field enabled
- Current interpolation across timestamp gaps greater than 36.0 h is outside-time support
- compatible=1660 marginal=56 rejected=3 missing_SST=41
- Isotope sensitivities: seawater SD=0.20?, calibration SD=1.00 deg C, residual SD=1.50 deg C correlated over 14.0 d, SST SD=0.50 deg C, shell smoothing=7.0 d, compatible/rejection simultaneous confidence=0.950/0.995, maximum marginal/rejection penalties=0.693/4.605 log units

Material limitations

- Diagnostic source-compatibility result, not a calibrated final MH370 posterior.
- Current families are reported separately and are never averaged using subjective weights.
- A2-G1 covers only the final months; the screen is neutral for any compatible chronology and never density-ranks compatible paths.
- No field-unavailable trajectory is relabelled as beaching; termination fractions and reasons remain in the JSON artifact.
- GLORYS12V1 is assimilative; the available drifter replay is not guaranteed held out.
- Daily means omit subdaily current variability represented by the native-HYCOM three-hour analyses.
- Land and any missing ocean value remain the packed missing sentinel; no missing value is converted to zero current.
- Stokes drift, direct windage, and coastal beaching are not included in this current field and require distinct explicit inputs.