

MH370 flaperon drift diagnostic - CMEMS GLORYS12 currents plus full WAVERYS surface Stokes

Uncalibrated family-specific diagnostic; currents=cmems_glorys12_reanalysis | isotope=enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival

Estimate at a glance

CURRENT FAMILY
cmems_glorys12_reanalysis

PEAK SOURCE CELL
arc-146

PEAK POSITION
-27.621 deg , 100.219 deg

WEIGHTED TRAJECTORIES
1024 x 77 cells


MEAN IMPORTANCE-WEIGHTED FIELD-TERMINATION FRACTION
46.5%

MEAN PROPOSAL TERMINATION FRACTION
26.8%

ARRIVAL SAMPLER
AdaptiveSmc

PEAK CORRECTION ESS
376.4


dark outline: 50% HDR
blue outline: 90% HDR



Probability is conditional on the named motion, SATCOM, and evidence configuration.

Posterior geography

Highest-density regions are computed from the weighted particle population.




Interpretation

The map is a probability density over the modelled impact state. It is not a searched-area elimination map, and structural model alternatives must be compared separately.

Numerical behaviour and conditions

Adaptive likelihood tempering, particle diversity, and declared evidence conditions.



Evidence conditions

- Current family selected explicitly: cmems_glorys12_reanalysis
- Current persistent-land corners renormalized=true; intermittent missing values remain fatal
- Barnacle isotope: enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival
- direct windage disabled
- distinct Stokes velocity field enabled at 100% surface response; finite coastal corners renormalized=true
- explicit land mask enabled; coastal beaching disabled because no beaching-rate field was supplied
- Current interpolation across timestamp gaps greater than 36.0 h is outside-time support
- compatible=711 marginal=12 rejected=1 missing_SST=200

Material limitations

- Diagnostic source-compatibility result, not a calibrated final MH370 posterior.
- Current families are reported separately and are never averaged using subjective weights.
- A2-G1 covers only the final months; the screen is neutral for any compatible chronology and never density-ranks compatible paths.
- No field-unavailable trajectory is relabelled as beaching; termination fractions and reasons remain in the JSON artifact.
- GLORYS12V1 is assimilative; the available drifter replay is not guaranteed held out.
- Daily means omit subdaily current variability represented by the native-HYCOM three-hour analyses.
- Land and any missing ocean value remain the packed missing sentinel; no missing value is converted to zero current.
- Stokes drift, direct windage, and coastal beaching are not included in this current field and require distinct explicit inputs.