


MH370 flaperon drift diagnostic - HYCOM family - oce

Diagnostic family-specific posterior; currents=hycom | isotope=enabled as P(isotope | represented flaperon arrival)

Estimate at a glance


CURRENT FAMILY	PEAK SOURCE CELL
hycom	arc-154
PEAK POSITION	PARTICLES
-31.683 deg , 96.747 deg	64 ? 77 cells
MEAN FIELD-TERMINATION FRACTION	
71.8%	



Probability is conditional on the named motion, SATCOM, and evidence configuration.

Posterior geography

Highest-density regions are computed from the weighted particle population.




Interpretation

The map is a probability density over the modelled impact state. It is not a searched-area elimination map, and structural model alternatives must be compared separately.

Numerical behaviour and conditions

Adaptive likelihood tempering, particle diversity, and declared evidence conditions.



Evidence conditions

- Current family selected explicitly: hycom
- Barnacle isotope: enabled as $P(\text{isotope} \mid \text{represented flaperon arrival})$
- direct windage disabled
- distinct Stokes velocity field enabled
- Recovery dates are discovery intervals with a configured pre-discovery encounter window; they are not asserted beaching times.

Material limitations

- Diagnostic source-area result, not a calibrated final MH370 posterior.
- The preserved HYCOM field is approximately 0.96 deg , not native 1/12 deg ; this can dominate long-trajectory error.
- GDP is a monthly climatology with incomplete ocean coverage and is not interchangeable with realized HYCOM currents.
- A2-G1 covers only the final months and its chronology is a leading uncertainty; the 49 target rows use a correlated residual model.
- No field-unavailable trajectory is relabelled as beaching; termination fractions and reasons remain in the JSON artifact.