

MH370 flaperon drift diagnostic - GDP family - ocean-

Uncalibrated family-specific diagnostic; currents=gdp | isotope=enabled as a neutral-or-penalizing screen conditional on repr

Estimate at a glance

CURRENT FAMILY

gdp

PEAK SOURCE CELL

arc-163

PEAK POSITION

-35.752 deg , 91.975 deg

WEIGHTED TRAJECTORIES

2048 ? 77 cells

MEAN IMPORTANCE-WEIGHTED FIELD-TERMINATION FRACTION

60.6%

MEAN PROPOSAL TERMINATION FRACTION

31.8%


ARRIVAL SAMPLER

AdaptiveSmc

PEAK CORRECTION ESS

870.5


dark outline: 50% HDR
blue outline: 90% HDR



Probability is conditional on the named motion, SATCOM, and evidence configuration.

Posterior geography

Highest-density regions are computed from the weighted particle population.




Interpretation

The map is a probability density over the modelled impact state. It is not a searched-area elimination map, and structural model alternatives must be compared separately.

Numerical behaviour and conditions

Adaptive likelihood tempering, particle diversity, and declared evidence conditions.



Evidence conditions

- Current family selected explicitly: gdp
- Barnacle isotope: enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival
- direct windage disabled
- distinct Stokes velocity field enabled
- coastal beaching disabled; no beaching rate was invented
- No additional current interpolation-gap cutoff configured
- compatible=1386 marginal=1 rejected=0 missing_SST=301
- Isotope sensitivities: seawater SD=0.20?, calibration SD=1.00 deg C, residual SD=1.50 deg C correlated over 14.0 d, SST SD=0.50 deg C, shell smoothing=7.0 d, compatible/rejection simultaneous confidence=0.950/0.995, maximum marginal/rejection penalties=0.693/4.605 log units

Material limitations

- Diagnostic source-compatibility result, not a calibrated final MH370 posterior.
- Current families are reported separately and are never averaged using subjective weights.
- A2-G1 covers only the final months; the screen is neutral for any compatible chronology and never density-ranks compatible paths.
- No field-unavailable trajectory is relabelled as beaching; termination fractions and reasons remain in the JSON artifact.
- This preserved subset is 0.5 degree; the current official product is 0.25 degree.
- Land and sparsely observed missing cells are not nearest-filled.
- Formal mean-current standard errors are not eddy diffusivity.