

MH370 stringent multi-debris GDP-climatology sensitivity - separated recovered and non-recovery populations

Uncalibrated family-specific diagnostic; currents=gdp | isotope=enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival

Estimate at a glance

CURRENT FAMILY

gdp

PEAK POSITION

-36.571 deg , 90.785 deg

RECOVERY EVIDENCE

9 objects in 9 independent episodes

MEAN PROPOSAL TERMINATION FRACTION

77.5%

PEAK SOURCE CELL

arc-165

WEIGHTED TRAJECTORIES

1024 x 77 cells x 11 transport ensembles (9 event-targeted n


MEAN IMPORTANCE-WEIGHTED FIELD-TERMINATION FRACTION

78.5%

ARRIVAL SAMPLER

AdaptiveSmc


dark outline: 50% HDR
blue outline: 90% HDR



Probability is conditional on the named motion, SATCOM, and evidence configuration.

Posterior geography

Highest-density regions are computed from the weighted particle population.




Interpretation

The map is a probability density over the modelled impact state. It is not a searched-area elimination map, and structural model alternatives must be compared separately.

Numerical behaviour and conditions

Adaptive likelihood tempering, particle diversity, and declared evidence conditions.



Evidence conditions

- Current family selected explicitly: gdp
- Object-motion families selected explicitly: measured-flaperon-response, low-exposure-exterior, uncertain-interior-panels-low-exposure-comparator, unobserved-low-exposure-non-recovery-population, unobserved-high-windage-non-recovery-population
- Current persistent-land corners renormalized=false; intermittent missing values remain fatal
- Barnacle isotope: enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival
- distinct direct 10 m windage: measured-flaperon-response=0.000% + 0.100 m/s at -20.0 degrees, unobserved-high-windage-non-recovery-population=3.000% + 0.000 m/s at 0.0 degrees
- distinct Stokes velocity field enabled; responses measured-flaperon-response=100%, low-exposure-exterior=100%, uncertain-interior-panels-low-exposure-comparator=100%, unobserved-low-exposure-non-recovery-population=100%, unobserved-high-windage-non-recovery-population=0%; finite coastal corners renormalized=true
- explicit land mask enabled; coastal beaching disabled because no beaching-rate field was supplied
- No additional current interpolation-gap cutoff configured

Material limitations

- Diagnostic source-compatibility result, not a calibrated final MH370 posterior.
- Current families are reported separately and are never averaged using subjective weights.
- A2-G1 covers only the final months; the screen is neutral for any compatible chronology and never density-ranks compatible paths.
- No field-unavailable trajectory is relabelled as beaching; termination fractions and reasons remain in the JSON artifact.
- This preserved subset is 0.5 degree; the current official product is 0.25 degree.
- Land and sparsely observed missing cells are not nearest-filled.
- Formal mean-current standard errors are not eddy diffusivity.
- This is the native wave model persistent support mask, not a measured shoreline-distance, beaching-probability, retention, or refloating model.