


# MH370 flaperon drift diagnostic - native-grid

## HYCOM currents only

Uncalibrated family-specific diagnostic; currents=hycom\_gofs31\_reanalysis | isotope=enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival


### Estimate at a glance

|                                                     |                                    |
|-----------------------------------------------------|------------------------------------|
| CURRENT FAMILY                                      | PEAK SOURCE CELL                   |
| hycom_gofs31_reanalysis                             | arc-121                            |
| PEAK POSITION                                       | WEIGHTED TRAJECTORIES              |
| -13.146 deg , 107.020 deg                           | 4096 x 77 cells                    |
| MEAN IMPORTANCE-WEIGHTED FIELD-TERMINATION FRACTION | MEAN PROPOSAL TERMINATION FRACTION |
| 9.6%                                                | 4.2%                               |
| ARRIVAL SAMPLER                                     | PEAK CORRECTION ESS                |
| AdaptiveSmc                                         | 1853.9                             |



# Posterior geography

Highest-density regions are computed from the weighted particle population.




## Interpretation

The map is a probability density over the modelled impact state. It is not a searched-area elimination map, and structural model alternatives must be compared separately.

# Numerical behaviour and conditions

Adaptive likelihood tempering, particle diversity, and declared evidence conditions.



## Evidence conditions

- Current family selected explicitly: hycom\_gofs31\_reanalysis
- Barnacle isotope: enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival
- direct windage disabled
- Stokes drift omitted
- explicit land mask enabled; coastal beaching disabled because no beaching-rate field was supplied
- Current interpolation across timestamp gaps greater than 36.0 h is outside-time support
- compatible=3350 marginal=231 rejected=8 missing\_SST=390
- Isotope sensitivities: seawater SD=0.20 per mil, calibration SD=1.00 deg C, residual SD=1.50 deg C correlated over 14.0 d, SST SD=0.50 deg C, shell smoothing=7.0 d, compatible/rejection simultaneous confidence=0.950/0.995, maximum marginal/rejection penalties=0.693/4.605 log units

## Material limitations

- Diagnostic source-compatibility result, not a calibrated final MH370 posterior.
- Current families are reported separately and are never averaged using subjective weights.
- A2-G1 covers only the final months; the screen is neutral for any compatible chronology and never density-ranks compatible paths.
- No field-unavailable trajectory is relabelled as beaching; termination fractions and reasons remain in the JSON artifact.
- This is every eighth available 3-hour analysis, not a daily mean; exact supplemental analyses repair two stride-created gaps, leaving variable 24-30 hour spacing and phase changes.
- The GOFS 3.1-like reanalysis documents known deep-water and Philippine Sea issues; predictive drifter validation is required before primary use.
- Land remains the packed missing sentinel. No missing ocean value is converted to zero current.
- Stokes drift, direct windage, and coastal beaching are not included in this current field and require distinct explicit inputs.