


MH370 flaperon drift diagnostic - native-grid

HYCOM reanalysis sensitivity

Uncalibrated family-specific diagnostic; currents=hycom_gofs31_reanalysis | isotope=enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival


Estimate at a glance

CURRENT FAMILY	PEAK SOURCE CELL
hycom_gofs31_reanalysis	arc-166
PEAK POSITION	WEIGHTED TRAJECTORIES
-36.968 deg , 90.173 deg	1024 x 77 cells x 1 transport ensembles (1 event-targeted reanalysis)
RECOVERY EVIDENCE	MEAN IMPORTANCE-WEIGHTED FIELD-TERMINATION FRACTION
1 objects in 1 independent episodes	73.3%
MEAN PROPOSAL TERMINATION FRACTION	ARRIVAL SAMPLER
69.0%	AdaptiveSmc



Posterior geography

Highest-density regions are computed from the weighted particle population.




Interpretation

The map is a probability density over the modelled impact state. It is not a searched-area elimination map, and structural model alternatives must be compared separately.

Numerical behaviour and conditions

Adaptive likelihood tempering, particle diversity, and declared evidence conditions.



Evidence conditions

- Current family selected explicitly: hycom_gofs31_reanalysis
- Object-motion families selected explicitly: configured-object-motion
- Current persistent-land corners renormalized=false; intermittent missing values remain fatal
- Barnacle isotope: enabled as a neutral-or-penalizing screen conditional on represented flaperon arrival
- distinct direct 10 m windage: configured-object-motion=0.000% + 0.100 m/s at -20.0 degrees
- distinct Stokes velocity field enabled; responses configured-object-motion=100%; finite coastal corners renormalized=false
- explicit land mask enabled; coastal beaching disabled because no beaching-rate field was supplied
- Current interpolation across timestamp gaps greater than 36.0 h is outside-time support

Material limitations

- Diagnostic source-compatibility result, not a calibrated final MH370 posterior.
- Current families are reported separately and are never averaged using subjective weights.
- A2-G1 covers only the final months; the screen is neutral for any compatible chronology and never density-ranks compatible paths.
- No field-unavailable trajectory is relabelled as beaching; termination fractions and reasons remain in the JSON artifact.
- This is every eighth available 3-hour analysis, not a daily mean; exact supplemental analyses repair two stride-created gaps, leaving variable 24-30 hour spacing and phase changes.
- The GOFS 3.1-like reanalysis documents known deep-water and Philippine Sea issues; predictive drifter validation is required before primary use.
- Land remains the packed missing sentinel. No missing ocean value is converted to zero current.
- Stokes drift, direct windage, and coastal beaching are not included in this current field and require distinct explicit inputs.