




Outliers from the H08S tests on Kadri (2024) Fig. 9 vectors (filtered, single channel; not raw IMS): cycle test h08s_shot_outliers.py (ddaa848/4b0ee7e), energy-only test h08s_pulse_energy.py (8ff37fb) and the exploratory off-cadence search. Each is paired with every H01W trigger (2a detector, strict or loose) or bearing-gated Kadri Table 1 event whose H08S–H01W time difference lies in 856–1,347 s. The implied source and impact time are prior-weighted means using both arrival times; impact 00:19:37 + 300 ± 180 s, $c = 1.482 \pm 0.006$ km/s, 10 s picks; Table 1 bearings with Student-t sd 3.3°. Filled = $\log_{10} \text{BF} \geq 1$ against a chance pair; open = below; bars = ± 1 sd. Every pre-registered count is consistent with chance ($p \geq 0.20$). The grey map prior and the grey impact-time prior are the stand-in from run no-exhaustion-prior (b3dd44b), prior track 295.66° ± 1.0°, config sensitivity/no-exhaustion-prior.toml, σ_x 20 NM. The blue curve is the reference-289 impact-time distribution from eof-289-full seed 1 (source run snap289-m0011, prior track 289.7° ± 1.0°, held-out 00:19 option, weight $\times \exp(\loglik:\text{none})$), shown for comparison only and not used in the scoring. EXPLORATORY characterisation.