With the present evidence the types of evidence do not converge: only the satellite data narrow the arc



Note (STE100): This chart measures how much each type of evidence narrows the part of the 7th arc where the aircraft can be. One bit halves the arc length. The satellite data and the flight model put the probability near 37 S. The debris drift and the searched areas, on their own, put more probability further north. So when we add them to the satellite result, the probable arc becomes longer, not shorter. The test in (d) shows that the satellite result and the other evidence disagree at about 2 sigma. This is a moderate disagreement, not proof of an error. The hydrophone data are not used. The Pleiades objects and the Australian coasts are separate cases, not the main result. Dots: 00:19 data held out. Grey ticks: the other five ways to use the 00:19 data.
Technical: composer views v3.2 machinery reused unchanged (compose_v3.py, V32=1 DEXT=1; weights reproduce products core, D, S, DS, ApopS, DSP, U-views with max relative difference 0). Core run J (tag 1244b7c), four-stratum mixture, settling run J (breakup table provisional), settled seabed; log-on cause competing, prior odds 1/3; end-of-flight uncontrolled arm not calibrated. Drift: finds only, arc-latitude profile 20-40 S, GLORYS12 + GlobCurrent marginalised jointly with the Australian and Pleiades terms (0.5 sum over models of the product). Searched areas: per-cell model V1 central, Ocean Infinity coverage inferred (grade C), included as best estimate. Australian non-finds: measured, populated coast only (labelled sensitivity). Hydroacoustics zero weight by ruling. Bits: entropy on 10 NM bins of the impact arc position against a uniform base on the arc where every module is computed (40.25-19.9 S; 28 % of the 40.6-11.8 S span is outside; core mass outside 0.015 %); views without core are u x E[L | bin] from core outcomes. (a) impact arc position; (b) arc length on 10 NM bins; (e) 2-D settled seabed on the 769 km2 equal-area grid. (c) Shapley values over all subsets of each set. (d) suspiciousness ln S = ln R - ln I with Bayesian model dimensionality (Handley and Lemos 2019), each type against the product of the others in its set; Gaussian calibration, approximate. Intervals: 400 seed-bootstrap draws (composer resampling groups), 95 %.