
If the imaged objects are from 9M-MRO: where the debris entered the sea — 00:19 R600 BTO Only
drift-error correlation 0.25, two ocean models (proposed headline) — core run C NOT CONVERGED



	ρ = 0.25 two models (drawn)	ρ = 0.25 GLORYS12 only	ρ = 0 two models	ρ = 0.5 two models
50 % region, km²	9,617	6,980	9,620	9,840
90 % region, km²	49,944 σ pending	34,997 σ pending	46,163 σ pending	53,468 σ pending
mean position	36.00 S 92.42 E	36.06 S 92.40 E	35.93 S 92.40 E	36.05 S 92.43 E
outside past searches	75.0 %	78.1 %	73.1 %	76.3 %
share left by searches	0.411	0.450	0.390	0.424
tension ln S (p)	-1.15 (0.16)	-0.67 (0.23)	-1.35 (0.14)	-1.00 (0.18)
mean shift from PDF without H	99 NM	95 NM	102 NM	96 NM
PDF without H inside 90 % region	8 %	7 %	7 %	9 %

- probability density if the objects are debris (relative)
- 50 % region (drawn setting)
- 90 % region (drawn setting)
- 90 % region, drift-error correlation 0.25, GLORYS12 only
- 90 % region, drift-error correlation 0, two ocean models
- 90 % region, drift-error correlation 0.5, two ocean models
- impact PDF without H (sampled points)
- ATSB Phase 2 2014-17 (official)
- Ocean Infinity 2018 (grade C)
- OI 2025-26 south-east band (grade C)
- OI north-west band (not searched; grade C)
- 7th arc, FL400
- Pléiades rating-5 objects
- COSMO-SkyMed radar contacts (all four)

STE100: This map shows where the aircraft possibly went into the sea, if the objects in the Pléiades satellite images are from 9M-MRO. The blue colour shows the probability with this assumption. The grey points show the probability without this assumption. The drawn result uses two ocean models with equal weight and a correlation of 0.25 between the drift errors of the two satellite data sets. The thin lines show the 90 % region for other settings. This is a proposed headline; the architecture session must accept it. The flight calculation (core run C) is not converged. Use these results with caution. Technical: core run C (4 strata x 8 seeds) split-half NOT CONVERGED (floor 0.924: free 0.918, Davey dynamics + radar 0.910, descent-climb 0.892, routes 0.850); end-of-flight physics PROVISIONAL (descents > 6,500 ft/min and unloading not reachable; no structural limit, G13). 00:19 R600 BTO Only, existence constraint 'unpowered'; strata re-weighted by the 00:19 evidence (family-evidence-run-c.json, key '00:19 R600 BTO Only +unpowered': next-c-free 0.590, next-c-repro-radar 0.239, next-c-descent-climb 0.145, next-c-routes 0.027). After Phase 2 + Bluefin-21 (official) + OI 2018 + OI 2025-26 SE band (grade C; NW band drawn only). Under H: 12 rating-5 Pléiades objects (6 clusters, 3 km, equal weight) x all four COSMO-SkyMed contacts, one debris field; bivariate-normal transport errors per component (prepare/rho_surface.py), ρ = 0.25 (ocean transport measured 0.07-0.41), ρ = 0 / 0.5 beside. GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight; GlobCurrent windage product-relative (drift audit F1: weights of c - dc, dc uniform 0.60-0.75 % U10 by 3-point quadrature, mass below 0 clamped to node 0; surfaces joint-surface-rho*-gcrel). PROPOSED headline (module, 11 Oct 18:19 UTC); in force until accepted: GLORYS12 only (ruling 2, 11 Oct 02:15 -0600), shown beside. Split-half σ for these settings is queued on the heavy-compute lock and is not shown yet. Conditional on H; no identity likelihood, no Bayes factor or P(H | data). Holland H1/H2 not estimable. Background: 6,000 points from the impact PDF without H after the same searches.