
If the imaged objects are from 9M-MRO: where the debris entered the sea, on GEBCO 2026 relief — 00:19 R600 BTO Only



STE100: This map shows where the aircraft possibly went into the sea, if the objects in the Pléiades satellite images are from 9M-MRO. The red bands show the 50, 90 and 99 % regions with this assumption. The points show the probability without this assumption. The drawn result uses two ocean models with equal weight and a correlation of 0.25 between the drift errors of the two satellite data sets. This is a proposed headline; the architecture session must accept it. The flight calculation (core run C) is not converged. Use these results with caution.

Technical: core run C (4 strata x 8 seeds) split-half NOT CONVERGED (floor 0.924: free 0.918, Davey dynamics + radar 0.910, descent-climb 0.892, routes 0.850); end-of-flight physics PROVISIONAL (descents > 6,500 ft/min and unloading not reachable; no structural limit, G13). 00:19 R600 BTO Only, existence constraint 'unpowered'; strata re-weighted by the 00:19 evidence (family-evidence-run-c.json, key '00:19 R600 BTO Only +unpowered': next-c-free 0.590, next-c-repro-radar 0.239, next-c-descent-climb 0.145, next-c-routes 0.027). After Phase 2 + Bluefin-21 (official) + OI 2018 + OI 2025-26 SE band (grade C; NW band drawn only). Under H: 12 rating-5 Pléiades objects (6 clusters, 3 km, equal weight) x all four COSMO-SkyMed contacts, one debris field; bivariate-normal transport errors per component (prepare/rho_surface.py), ρ = 0.25 (ocean transport measured 0.07-0.41), ρ = 0 / 0.5 beside. GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight; GlobCurrent windage product-relative (drift audit F1: weights of c - dc, dc uniform 0.60-0.75 % U10 by 3-point quadrature, mass below 0 clamped to node 0; surfaces joint-surface-rho*-gcrel). PROPOSED headline (module, 11 Oct 18:19 UTC); in force until accepted: GLORYS12 only (ruling 2, 11 Oct 02:15 -0600), shown beside. Split-half σ for these settings is queued on the heavy-compute lock and is not shown yet. Conditional on H; no identity likelihood, no Bayes factor or P(H | data). Holland H1/H2 not estimable.

Result, after all searches: 90 % region 49,944 km², mean 36.00 S 92.42 E; tension ln S -1.15 (p 0.16), mean shift 99 NM, PDF without H inside the 90 % region 8 %. Beside it, GLORYS12 only at ρ = 0.25: 34,997 km², 36.06 S 92.40 E, ln S -0.67.

Relief: GEBCO 2026 15-arcsecond grid (doi:10.5285/4f68d5c7-45eb-f999-e063-7086abc036fa), shown at 0.025 deg; contours every 1 km depth. Right: probability per degree of latitude over 85-103 E, after all searches.

Background: the impact PDF without H (flight posterior alone, after the same searches), as 6,000 points drawn in proportion to its probability from 0.05 deg grid cells and jittered within each cell. 31 % of it lies inside this frame.