
Each added 00:19 measurement narrows the impact region; the R1200 frequency moves its peak to 37.7° S



Note 1: Each panel shows the area that holds 50, 90 and 99 % of the probability of the ocean impact, for one way to use the two 00:19 satellite messages. The heavy black, brown and green lines show sea floor that was searched; the search result is not applied in these maps. The log-on cause mixes fuel exhaustion (weight 3) and other causes (weight 1). The results are provisional.
Note 2: End of flight on core run I (4 route families x 8 seeds x 100,000 hand-off particles at 00:11; N = 8 children x 4 descents; family weights 0.9982 / 0.0011 / 0.0004 / 0.0002; binary mh370-eof-ri1). Competing causes C1 (Erlang lag) + C2 (1/W, W = 00:10:58-01:15:56 UTC, silent), q = 1/3; survival factor applied; no treatment uses the R1200 BTO. Mixed = equal-prior average of D3/D4/D5 weighted by ln Z (they score the same three observations). Density: 0.02° grid, Gaussian sigma 0.08°. Searched ground: coverage >= 0.5 outline of the seabed-search rasters (Bluefin-21 2014 not drawn). PROVISIONAL: piloted-turn model, INTERIM pitch-down, SLS thrust 415.45 kN (pre-correction, +2.1 %). Core run I not converged (8-seed floor 0.916). Effective impacts below 1,000 in D3/D5 x C2-silent cells (COVERAGE, eof-run-i-oct12). Enclosed-mass check (held out): 50.0 / 90.0 / 99.0 %.