Seabed wreckage latitude after fuel exhaustion with no control (A1):
the extra A1 run reduces the sampling noise and does not move the peak



(i) The lines show where the wreckage is on the seabed, by latitude, for the flight end 'cruise to fuel exhaustion, then no control' (A1). Each panel uses one way to read the 00:19 satellite data. The dashed grey line uses only the main flight-end run. The blue line also uses an extra run that samples only this flight end; a known scale factor keeps the probabilities correct. The thin blue lines show the two halves of the extra run (each with the main-run part outside the extra run). Some parts of the flight model are provisional. Search results and debris drift are not used.

(ii) Settled-mass-weighted seabed latitude of A1 outcomes (afloat pieces excluded), 0.1° bins, systematic resample N = 10,000, one settling draw each; settling 9330d4c (GLORYS12V1, TEOS-10, ERA5, AusSeabed/GEBCO_2026; breakup table PROVISIONAL; breakup rows from the intact impact point, G-bf). Prior track: core run J waypoint routes (stratum conditional, core share 0.9982). Impacts: end of flight run J (24 seeds, mh370-eof-rj1) and the A1-branch conditional pass (8 seeds, prior on onset flame-out-associated + no intervention, D3-aimed scored selection, exact correction) combined by the stratified estimator, P_D0'(branch) = 0.0935. Primary cell: both log-on causes, q = 1/3, C2 silent; Held Out is D0' (impact after 00:19:37.443). Uncontrolled-arm dynamics not calibrated (Boeing leave-one-out gate failed). Row ESS by Kish, gate 1,000: the main run under both frequencies as recorded (368) is below it. Core run J split-half accepted as a pass by project ruling, 16 Oct 2026. Davey et al. (2016) Fig. 10.3 not drawn: it is a flight posterior, not an A1-conditional seabed density.