


If the imaged objects are from the aircraft (H): where the aircraft hit the sea, with the tension against the flight data alone — UNCONVERGED



(i) The blue shading shows where the aircraft hit the sea if the floating objects seen by the satellites came from it. The grey points show the same without that assumption. The searches of the seabed are not applied in this chart. Panel a uses the 00:19 range and frequency values; panel b uses only the 00:19 range value. The objects move the estimate 100 to 135 NM (mean) north-east, into a small part of the area that the flight data alone allow. The flight model is not stable yet (it has not converged), so all values can change. This work was done by a stand-in for the Pléiades team, which must review it. The ± values are the differences between two halves of the runs.

(ii) $p(x | D, H) = \text{mixture over core (b) families of the end-of-flight + alive impact posterior } x | L_H, L_H = \text{Pléiades rating-5 (6 clusters, equal weights) + COSMO-SkyMed C4 pass-marginal, GLORYS12+ERA5 and GlobCurrent daily+ERA5 at equal weight, transport errors independent (rho = 0; rho = 0.5 not supported by the hook, see note), GlobCurrent windage as run (drift audit F1), debris released 00:20 UTC; surfaces sha256 9a3d55a9/e7fd6ded (hypothesis/pleiades 74332d0). Impacts mh370-exchange/end-of-flight/next-run (4 strata x 4 seeds, 51.2 M rows; EoF physics provisional; two-tank bookkeeping only); log-on cause other. Strata mixed by end of flight's 00:19-re-weighted P(family) (family-evidence-next-run-b.json; ruling C), fixed P(family) beside. 0.05 deg grid 85-103 E, 43-25 S (in-grid mass without H 0.994). Tension (rerun_reference.tension): ln R = ln[E_flight(L_H) / grid-mean(L_H)]; HDR overlaps two-way; mode on the unsmoothed grid (noisy without H). ± = split-half sigma, rms over the three 2+2 seed partitions of (half A - half B)/2. ESS = Kish, mixture weights. Holland H1/H2 not estimable. Searches not applied. Stand-in run 10 Oct 2026, module to review.$