




(i) Each line is one flight path drawn at random in proportion to its probability (about 240 per panel). Grey: the flight data alone. Blue: if the floating objects came from the aircraft. From 00:11 to the sea, each blue line is the same calculated path that the objects support. Before 00:11 the files keep no link from a path to its later part, so the light-blue lines are the general route sample, given more weight only for the autopilot modes that the objects support. They are not the exact earlier part of the blue lines. Under H the paths at 00:11 are further north (34-35 S), slower and turned less to the west. The flight model has not converged.

(ii) Option r600/no-offset+alive (log-on cause other); strata mixed by EoF's 00:19-re-weighted P(family), under H by $P(f) Z_{f(H)}/\text{sum}$. 00:11-to-impact lines: Efraimidis-Spirakis weighted draws without replacement (80 per seed and case, weights $w \propto \exp(l_i) \times \text{alive}$, and $x_{L,H}$ under H), n_f per stratum in proportion to the stratum share; vertices hand-off m0011 (handoff.npy via 'parent'), EoF take-over, position at 00:19:37 (latent last burst), impact, in time order. 18:01-00:11 lines: core routes.npy (2,000 per seed, 600 s, final-posterior mode sample, NOT linked to hand-off rows; final_row = i64::MAX at m0011); mode of each route assigned from diagnostics.json posterior_probability boundaries (+/-1 route per boundary); under H re-weighted by that stratum's hand-off share ratio H/no-H for the filter mode. L_H as Fig. 1 (P + C4, two ocean models, $\rho = 0$). Core (b) split-half NOT converged; EoF physics provisional.