




(i) A point to the right of the dotted line means that this kind of flight path becomes more probable if the floating objects came from the aircraft. A point to the left means less probable. The bars show the run-to-run spread. Rows for 00:11 and earlier come from the flight model; rows for the descent come from the end-of-flight model. The flight model has not converged, so treat the values as provisional. A stand-in made this chart for the Pléiades team, which must review it.

(ii) Shares are posterior shares of the mixture over core (b) strata (P(family) re-weighted by the 00:19 data, ruling C), without H (EoF arm weights $w \times \exp(\|l\| \times \text{alive}, \log\text{-on cause other})$ and under H (x L H: Pléiades rating-5 + COSMO C4, two ocean models, $\rho = 0$, GlobCurrent windage as run). 00:11 and earlier: aggregated over impacts to hand-off parents ('parent' -> handoff-m0011 rows: handoff.npy + handoff.toml state). Descent rows: end-of-flight latents per impact. sigma = split-half, rms over the three 2+2 seed partitions of (half A - half B)/2; bars +/- 2 sigma. ESS = Kish under H per category (paths: hand-off parents; descent: impacts). Bins with share < 0.5 % without H omitted from the chart (all bins in the CSV). EoF physics provisional; known coverage gaps (architecture list): EoF commanded descent rates capped at 6,500 ft/min, free flight cannot unload, family B onset; Holland H1/H2 not estimable.