




(i) The searches of the seabed found nothing. If the floating objects came from the aircraft, the searches remove about 64 % of the probability, against about 49 % without that assumption, because the objects point to places near searched ground. The blue area that remains is next to the searched strips. The value of ρ (the chance that the sonar missed the wreck) changes the removed share most. Ocean Infinity areas are traced from public sources (grade C). The flight model has not converged. A stand-in made this chart; the teams must review it.

(ii) Searched areas' likelihood, its own code: $P(\text{no find}) = \rho + (1-\rho) \text{prod}_k(1 - q_k f_k c_k)$, shared misses, point target; c_k = covered_fraction_campaign from `mh370 evaluate hypotheses/seabed-search/run.toml` + OI 2018 and 2025 campaign lists` (binary at e9cca3b; base loglik reproduced to 0.0); Phase 2 q 0.945, Bluefin-21 q 0.9, OI 2018 q 0.9 f 0.889, OI 2025-26 SE band q 0.9 f 0.7808. Removed = 1 - Z_search; in-grid (85-103 E, 43-25 S) values in the table, whole-posterior values in panel c and the CSV (without H whole: 0.494). H as Fig. 1 (P + C4, two ocean models, rho_transport = 0). Strata re-weighted by 00:19 P(family). sigma = split-half over 2+2 seed partitions. Other sensitivities (q 0.90/0.98, independent misses, OI 2018 coverage 0.952) in the note. Core (b) NOT converged; EoF provisional.