

Question 2:

Let Y_1 have a $\text{Poisson}(\lambda_1)$ distribution and let Y_2 be independently distributed as a $\text{Poisson}(\lambda_2)$.

Derive the probability mass function of $Y = Y_1 + Y_2$.

$$P_{Y_1}(y_1) = e^{-\lambda_1} \frac{\lambda_1^{y_1}}{y_1!}$$

$$P_{Y_2}(y_2) = e^{-\lambda_2} \frac{\lambda_2^{y_2}}{y_2!}$$

$$Y = Y_1 + Y_2$$

$$P_Y(t) = \sum_{y_1=0}^t e^{-\lambda_1} \frac{\lambda_1^{y_1}}{y_1!} e^{-\lambda_2} \frac{\lambda_2^{t-y_1}}{(t-y_1)!}$$

$$= e^{-\lambda_1 - \lambda_2} \frac{(\lambda_1 + \lambda_2)^t}{t!} \sum_{y_1=0}^t \frac{1}{y_1!} \frac{t!}{(t-y_1)!} \frac{\lambda_1^{y_1}}{(\lambda_1 + \lambda_2)^{y_1}} \frac{\lambda_2^{t-y_1}}{(\lambda_1 + \lambda_2)^{t-y_1}}$$

$$= \left(e^{-(\lambda_1 + \lambda_2)} \frac{(\lambda_1 + \lambda_2)^t}{t!} \right) \times 1, \text{ sum of binomial prob.}$$

$$\text{Poisson}(\lambda_1 + \lambda_2)$$

$$Y_1 \sim \text{Bin}(t, \frac{\lambda_1}{\lambda_1 + \lambda_2})$$

$$\text{Bin}(n, p)$$

$$\left(\frac{n!}{y!(n-y)!} \right) p^y (1-p)^{n-y}$$

$$\sum_{y=0}^n$$

$$= 1$$

Question 3:

Let X_1, X_2, \dots, X_n be independent $\text{Exponential}(\lambda)$ random variables. Derive the CDF and the PDF of $X_{(n)}$ where $X_{(n)} = \max_i X_i$.

$$F_{X_{(n)}}(x) = P(X_{(n)} \leq x)$$

$$\max_i X_i \leq x \text{ i.e.}$$

$$X_1 \leq x \text{ and } X_2 \leq x \dots \text{and } X_n \leq x$$

$$P(\max_i X_i \leq x) = P(X_1 \leq x) \times P(X_2 \leq x) \times \dots \times P(X_n \leq x)$$

$$= \prod_{i=1}^n F_{X_i}(x)$$

CDF for $\text{Exponential}(\lambda)$

$$1 - e^{-\lambda x} \quad \begin{matrix} x > 0 \\ \lambda > 0 \end{matrix}$$


$$P(\max X_i \leq x) = \prod_{i=1}^n (1 - e^{-\lambda x})$$

$$F_{X_{(n)}}(x) = (1 - e^{-\lambda x})^n$$

PDF $\frac{d}{dx} F_{X_{(n)}}(x) = n(1 - e^{-\lambda x})^{n-1} (-1) e^{-\lambda x} (-\lambda)$

$$= n(1 - e^{-\lambda x})^{n-1} \lambda e^{-\lambda x}$$

$$\max_i X_i \leq x$$



$$\max_i X_i \leq x \text{ iff all } X_i\text{'s} \leq x$$

$$= X_1 \leq x \text{ and } X_2 \leq x \dots X_n \leq x$$

$$\begin{aligned}
 & P_n(X_1 \leq x \cap X_2 \leq x \cap \dots \cap X_n \leq x) \\
 &= \underline{P_n(X_1 \leq x)} \times \dots \times \underline{P(X_n \leq x)} \\
 &= (1 - e^{-\lambda x})^n
 \end{aligned}$$

$\min X_i \leq x$ iff at least one $X_i \leq x$

$1 - F(x)$ $(\min X_i > x)$ iff

$$\begin{aligned}
 & \downarrow \quad X_1 > x \cap X_2 > x \cap \dots \cap X_n > x \\
 & P(\quad) = (P(X_i > x))^n
 \end{aligned}$$

Question 1:

A random rectangle has sides whose lengths are independent random variables, each with a uniform distribution on the interval $(0, 2)$. Find the expected value of the area and the expected value of the perimeter of the square.


$$X, Y \sim \text{ind } U(0, 2)$$

$$E(XY) = E(X)E(Y)$$

$$E(X) = 1 = E(Y)$$

$$\Rightarrow 1 \cdot 1 = 1$$

$$E(2X + 2Y) = 4$$



Chapter 4.5 Moment-generating functions

Distributions for random variables can be represented in many ways. We've seen some:

CDF: $F(x) = P(X \leq x)$ ✓ ✓ ✓

pmf: $p(x) = P(X = x)$

pdf: $f(x) : P(X \in A) = \int_A f(x) dx$

We know that $E(X)$ does not determine a distribution unless restricting to a special family: e.g. Poisson or exponential.


$$N(\mu, \sigma^2)$$

The moment-generating function (MGF) is another way that works for many distributions.

Suppose a function has a Taylor series expansion around 0, (Maclaurin series)

$$\underline{f(t)} = \underline{a_0} + \underline{a_1 t} + \underline{a_2 t^2} + \underline{a_3 \frac{t^3}{3!}} + \underline{a_4 \frac{t^4}{4!}} + \underline{a_5 \frac{t^5}{5!}} + \dots + \underline{a_n \frac{t^n}{n!}} + \dots$$

and the power series converges in an open interval around 0.



Then

1) $\frac{d^n}{dt^n} f(0) = a_n$ and has derivatives of all orders.

2) So the derivatives of f at 0
determine the Taylor series in radius of convergence.
and thus determine the entire function.

Defn: The n th moment of X is $E(X^n)$ if it exists.
central $E[(X - E(X))^n]$

Note: $\text{Var}(X)$ is the second central moment.

4
puntos

$E[(X - \mu_x)^3]$ is the "skewness"
sometimes standardized as $E\left[\left(\frac{X - \mu_x}{\sigma_x}\right)^3\right]$

MGF for r.v. X :

$$C(t) = E(e^{itx})$$

A function of t defined as:

$$M(t) = E(e^{tx}) = \begin{cases} \int e^{tx} f(x) dx \\ \sum_x e^{tx} p(x) \end{cases}$$

provided $|m(t)| < \infty$ in an interval $(-\varepsilon, \varepsilon)$, $\varepsilon > 0$.

$$M'(t) = \frac{d}{dt} \int_{-\infty}^{\infty} e^{tx} f(x) dx$$

$$= \int_{-\infty}^{\infty} \left(\frac{d}{dt} e^{tx} \right) f(x) dx$$

$$= \int_{-\infty}^{\infty} x e^{tx} f(x) dx$$

So $\Lambda'(0) = \int_{-\infty}^{\infty} x e^{0x} f(x) dx = E(X)$

$$\Lambda''(t) = \int_{-\infty}^{\infty} x^2 e^{tx} f(x) dx$$

$$\Lambda''(0) = E(X^2)$$

$$\Lambda^{(k)}(0) = E(X^k)$$

Two theorems:

If $M_X(t)$ exists in an open interval about 0 then

1) $E(X^k) = \Lambda^{(k)}(0)$

2) The distribution of X is uniquely determined by Λ

So the MGF joins the CDF, pmf and pdf as ways of characterizing distributions.

Note: All distribution for RVs
have a (unique) CDF.

Only some have a pdf, pmf or mgf.
- Both continuous and discrete
distribution can have mgf's.

Some mgf's

Poisson: $P(x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad x=0, 1, \dots; \lambda > 0$

$$\Lambda(t) = \sum_{x=0}^{\infty} e^{tx} \frac{\lambda^x}{x!} e^{-\lambda} \quad \sum x^n \frac{\lambda^x}{x!} e^{-\lambda}$$

$$= e^{-\lambda} \sum_{x=0}^{\infty} \frac{(\lambda e^t)^x}{x!} = e^{-\lambda} e^{\lambda e^t}$$

$$= e^{\lambda(e^t - 1)} \quad \text{Converges for all } t$$

Differentiating to find moments is much easier than summing infinite series!

$$M'(t) = e^{\lambda(e^t - 1)} \times \lambda e^t$$

$$M'(0) = e^{\lambda(e^0 - 1)} \times \lambda e^0 = e^0 \times \lambda e^0 = \lambda$$

$$M''(t) = \underbrace{e^{\lambda(e^t - 1)} \times \lambda e^t}_{\lambda} \times \underbrace{\lambda e^t}_{\lambda} + \underbrace{e^{\lambda(e^t - 1)}}_1 \underbrace{\lambda e^t}_{\lambda}$$

$$M''(0) = \lambda \quad \lambda + 1 \quad \lambda = \lambda^2 + \lambda$$

$$\begin{aligned}
 \text{So } \text{Var}(X) &= E(X^2) - (E(X))^2 \\
 &= (\lambda^2 + \lambda) - \lambda^2 \\
 &= \lambda
 \end{aligned}$$

What happens if we add 2 independent Poisson(λ): Y_1 and Y_2

Finding the pdf / pmf of a sum was tough. Let's see the MGF.

$$\begin{aligned}
 E(e^{t(Y_1 + Y_2)}) &= E(e^{tY_1})E(e^{tY_2}) \text{ by independence} \\
 &= e^{\lambda(e^t - 1)} e^{\lambda(e^t - 1)}
 \end{aligned}$$

$$M_{Y_1+Y_2}(t) = e^{2\lambda(e^t-1)}$$

Recognize this?

It's the MGF for a Poisson (2λ)

∴ by uniqueness of MGF

$$Y_1 + Y_2 \sim \text{Poisson}(2\lambda)$$

How about $Y_1 + Y_2 + \dots + Y_n$

where Y_i 's are independent

Poisson $\lambda_1, \lambda_2, \dots, \lambda_n$?

$$Y_1 + Y_2 + \dots + Y_n \sim \text{Poisson}(\sum \lambda_i)$$

Facts about MGFs

If the mgf exists in an interval around 0 then

1) $M^{(r)}(0) = E(X^r)$

2) If X_1, X_2, \dots, X_n are independent with mgfs $M_1(t), \dots, M_n(t)$, then

$$M_{X_1 + \dots + X_n}(t) = \prod_{i=1}^n M_i(t)$$

3) ... with the same mgf $M(t)$ then

$$M_{X_1 + \dots + X_n}(t) = [M(t)]^n$$

4) If $Y = a + bX$ then $M_Y(t) = e^{at} M_X(bt)$

Some LMGFs :

Exponential (λ)

$$\begin{aligned} \Lambda_{\lambda}(t) &= \int_0^{\infty} e^{tx} \lambda e^{-\lambda x} dx \\ &= \lambda \int_0^{\infty} e^{-(\lambda-t)x} dx \\ &= \frac{\lambda}{\lambda-t} \int_0^{\infty} (\lambda-t) e^{-(\lambda-t)x} dx \\ &= \frac{\lambda}{\lambda-t} \quad \text{if } \lambda-t > 0 \\ &\quad \text{i.e. } t < \lambda \end{aligned}$$

$$\text{Gamma}(\underset{\substack{\uparrow \\ \text{shape}}}{\alpha}, \lambda) = \left(\frac{\lambda}{\lambda - t} \right)^{\alpha}$$

So sum of n independent Exponential (λ)'s

$$\text{Gamma}(n, \lambda)$$

$$\text{Normal}(0, 1) \quad e^{-t^2/2}$$

Normal mean μ and variance σ^2

$$Y = \mu + \sigma Z \quad \text{where } Z \text{ is } N(0, 1)$$

$$\Lambda_{\mathbf{y}}(t) = E(e^{t \cdot \mathbf{y}}) = E(e^{t(\mu + \sigma Z)})$$

$$= E(e^{t\mu + t\sigma Z})$$

$$= e^{t\mu} E(t\sigma Z)$$

$$= e^{t\mu} e^{(t\sigma)^2/2}$$

$$= e^{t\mu + t^2\sigma^2/2}$$

Does it work?

$$\lambda'_y(t) = e^{t\mu + \frac{t^2}{2}\sigma^2} (\mu + t\sigma^2)$$

$$\begin{aligned} \lambda\lambda''_y(t) &= e^{t\mu + \frac{t^2}{2}\sigma^2} (\mu + t\sigma^2) (\mu + t\sigma^2) \\ &\quad + e^{t\mu + \frac{t^2}{2}\sigma^2} \sigma^2 \end{aligned}$$

$$= e^{t\mu + \frac{t^2}{2}\sigma^2} \{ (\mu + t\sigma^2)^2 + \sigma^2 \}$$

$$\text{So } E(Y) = M'_Y(0) = e^0 (\mu + 0 \cdot \sigma^2) = \mu$$

$$E(Y^2) = M''_Y(0) = e^0 \{ \mu^2 + \sigma^2 \} = \mu^2 + \sigma^2$$

$$\begin{aligned} \text{Var}(Y) &= E(Y^2) - (E(Y))^2 \\ &= \mu^2 + \sigma^2 - \mu^2 = \sigma^2 \end{aligned}$$

Sums of independent Normals

$$Y_1 \sim N(\mu_1, \sigma_1^2) \quad , \quad Y_2 \sim N(\mu_2, \sigma_2^2)$$

$$E(e^{t(Y_1+Y_2)}) = e^{t\mu_1 + \frac{t^2}{2}\sigma_1^2 + t\mu_2 + \frac{t^2}{2}\sigma_2^2}$$

$$= \exp\left\{t(\mu_1 + \mu_2) + \frac{t^2}{2}(\sigma_1^2 + \sigma_2^2)\right\}$$

$$\therefore Y_1 + Y_2 \sim N(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2) \quad !!!$$

